

PRODUCT MONOGRAPH

MENJUGATE[®] Liquid

10 micrograms suspension for injection

Meningococcal Group C Conjugate Vaccine

House Standard

Suspension for Injection

Active Immunizing Agent
ATC Code J07A H

GlaxoSmithKline Inc.
7333 Mississauga Road
Mississauga, Ontario
L5N 6L4

Date of Preparation:
September 15, 2015

Date of Approval: December 16, 2015

Control #: 189103

©2015 GlaxoSmithKline Inc., All Rights Reserved

Menjugate[®] is a registered trademark of GlaxoSmithKline Biologicals SA.

Table of Contents

PART I: HEALTH PROFESSIONAL INFORMATION.....	3
SUMMARY PRODUCT INFORMATION	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS	3
WARNINGS AND PRECAUTIONS.....	3
ADVERSE REACTIONS.....	5
DRUG INTERACTIONS	9
DOSAGE AND ADMINISTRATION.....	10
OVERDOSAGE	11
ACTION AND CLINICAL PHARMACOLOGY	11
STORAGE AND STABILITY.....	12
DOSAGE FORMS, COMPOSITION AND PACKAGING	12
PART II: SCIENTIFIC INFORMATION	14
PHARMACEUTICAL INFORMATION.....	14
CLINICAL TRIALS.....	14
DETAILED PHARMACOLOGY	16
TOXICOLOGY	16
PART III: CONSUMER INFORMATION.....	20

MENJUGATE® Liquid

Meningococcal Group C Conjugate Vaccine

PART I: HEALTH PROFESSIONAL INFORMATION

SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength per 0.5mL dose of vaccine	Clinically Relevant Nonmedicinal Ingredients
Intramuscular (IM) Injection	Suspension for injection 10 µg Meningococcal Group C Oligosaccharide conjugated to 12.5 to 25 µg CRM197	Aluminum hydroxide adjuvant <i>For a complete listing see Dosage Forms, Composition and Packaging section.</i>

INDICATIONS AND CLINICAL USE

Menjugate is indicated for the active immunization of children from 2 months of age, adolescents and adults, for the prevention of invasive disease caused by *Neisseria meningitidis* serogroup C.

CONTRAINDICATIONS

Menjugate is contraindicated in persons with a known hypersensitivity to any component of the vaccine and in persons who have shown signs of hypersensitivity after previous administration of Menjugate.

As with other vaccines, administration of Menjugate should be postponed in subjects with an acute severe febrile illness.

WARNINGS AND PRECAUTIONS

General

Before the injection of any biological, the health professional responsible for administration should take all precautions known for the prevention of allergic or any other reactions. As with all injectable vaccines, appropriate medical treatment and supervision should always be readily available in case of a rare anaphylactic event following administration of the vaccine.

Prior to administration of any dose of Menjugate (Meningococcal Group C-CRM 197 Conjugate Vaccine), the vaccine recipient (or parent or guardian) should be asked about personal history, family history, and recent health status, including immunization history, current health status and any adverse event associated with previous immunizations.

Menjugate will not protect against meningococcal diseases caused by any of the other types of

meningococcal bacteria (A, B, 29-E, H, I, K, L, W-135, X, Y, or Z, including non-typed). Complete protection against meningococcal serogroup C infection cannot be guaranteed.

Although symptoms of meningism such as neck pain/stiffness or photophobia have been reported, there is no evidence that the vaccine causes meningococcal C meningitis. Clinical alertness to the possibility of co-incidental meningitis should therefore be maintained.

Conjugate vaccines containing Cross Reacting Material 197 (CRM197) should not be considered as immunizing agents against diphtheria. No changes in the schedule for administering vaccines containing Diphtheria Toxoid are recommended.

Any acute infection or febrile illness is reason for delaying the use of Menjugate except when, in the opinion of the physician, withholding the vaccine entails a greater risk. A minor afebrile illness, such as a mild upper respiratory infection, is not usually reason to defer immunization.

The vaccine must not be injected intravenously, subcutaneously or intradermally.

Parents should be informed of the immunization schedule for this vaccine. Precautions such as use of antipyretic measures should be relayed to the parent or guardian, as well as the need to report any adverse event.

Latex-sensitive individuals - for syringe presentation:

Although no natural rubber latex is detected in the syringe tip cap, the safe use of Menjugate in latex-sensitive individuals has not been established.

Hematologic

Menjugate has not been evaluated in persons with thrombocytopenia or other bleeding disorders. The risk versus benefit for persons at risk of hemorrhage following intramuscular injection must be evaluated.

Immune

In individuals deficient in antibody production, vaccination may not result in an appropriate protective antibody response. While HIV infection is not a contraindication to vaccination, Menjugate has not been specifically evaluated in an immunocompromised population. Individuals with complement deficiencies and individuals with functional or anatomical asplenia may mount an immune response to meningococcal C conjugate vaccines; however, the degree of protection that would be afforded is unknown.

Special Populations

Pregnant Women: Animal studies have not demonstrated a risk to the fetus following administration of Menjugate. However, since no specific studies in humans have been carried out, caution is advised. The vaccine should not be used during pregnancy unless there is defined risk of meningococcal C disease, in which case the risk-benefit ratio should be evaluated.

Nursing Women: The effect on breast-fed infants of the administration of Menjugate to their

mothers has not been studied. The risk-benefit ratio should be examined before making the decision as to whether to immunize during lactation.

Geriatrics (> 65 years of age): There are no data in adults aged 65 years and older.

ADVERSE REACTIONS

Clinical Trial Adverse Drug Reactions

Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in the clinical trials may not reflect the rates observed in practice and should not be compared to the rates in the clinical trials of another drug. Adverse drug reaction information from clinical trials is useful for identifying drug-related adverse events and for approximating rates.

In controlled clinical studies performed in all age groups, signs and symptoms were actively monitored and recorded on diary cards following administration of the vaccine.

Of the local solicited symptoms, the most frequently reported were injection-site pain, erythema and swelling, which were normally mild and resolved within 24-72 hours following vaccination.

The general symptoms that have been solicited and reported were predominantly mild and resolved spontaneously. These include headache, malaise, nausea, arthralgia and myalgia in adolescents and adults; and irritability, change in appetite, diarrhea and fever in younger children. These solicited general symptoms were also reported in the control groups and have been reported when Menjugate (Meningococcal Group C–CRM197 Conjugate Vaccine) was administered concomitantly with other vaccines.

In infants and children aged 12 through 23 months, symptoms including crying, irritability, drowsiness, impaired sleeping, anorexia, diarrhea and vomiting were common after vaccination but there was no evidence that these were related to Menjugate rather than concomitant vaccines, particularly DTP.

The safety of Menjugate liquid formulation was compared with that of Menjugate lyophilised formulation in a randomised clinical study involving 989 children aged 12 months to 2 years. The safety profile of both formulations of Menjugate was comparable.

Children (12 through 23 months of age) Through Adults:

Table 1 presents an analysis of local and systemic reactions occurring within 7 days after one immunization with Menjugate. Data are pooled from 11 studies, representing approximately 1400 subjects. Most local and systemic reactions occurred by day 1 following immunization. In general, lower percentages of local and systemic reactions were present on days 2 through 6 following the first immunization.

Table 1 - Summary of Local and Systemic Post immunization Reactions Within 7 Days Following One Immunization of Menjugate, by Age Group at Enrollment*

	Percentage of Subjects		
	Menjugate 1-2 years n=942 (%)	Menjugate 3-5 years n=198 (%)	Menjugate 11-64 years n=269 (%)
Injection Site			
Pain (Any)	22	25	81
Severe	<1	0	2
Temperature (Any)	15	5	47
Hot	<1	1	8
Erythema (Any)	28	16	19
>50 mm	<1	0	1
Induration (Any)	16	7	24
>50 mm	<1	0	1
Systemic			
Change in Eating Habits	16	6	-
Sleepiness	19	9	-
Unusual Crying	4	1	-
Persistent Crying	1	0	-
Irritability	30	10	-
Vomiting	9	5	-
Diarrhea	18	8	-
Rash	9	4	-
Chills	-	-	13
Nausea	-	-	16
Malaise	-	-	25
Myalgia	-	-	29
Arthralgia	-	-	16
Headache	-	-	34
Temp $\geq 38^{\circ}\text{C}$	9	4	2
Stayed Home Due to Reaction	-	-	7
Analgesic/ Antipyretic Medication used	25	9	18

*This is a summary of data derived from a meta-analysis of 11 studies conducted in the United States, United Kingdom, Netherlands, and Canada. The recording of systemic reactions varied by age group, not all reactions were collected in all studies.

In clinical studies where subjects received Menjugate or a meningococcal polysaccharide vaccine, the rates of local pain and warmth were significantly lower with Menjugate in children 1 to 2 and 3 to 5 years of age; no differences were seen in the older subjects. In children 3 to 5 years of age, severe pain was seen in 9% of subjects with the polysaccharide vaccine and no subjects with Menjugate. The systemic reactions that were significantly less common in Menjugate subjects were fever, change in eating habits, irritability, and analgesic/antipyretic use in children aged 12 through 23 months of age, and irritability and analgesic/antipyretic use in children 3 to 5 years of age.

In adolescents and adults, the rates of all postimmunization reactions were similar after

Menjugate or polysaccharide vaccine administration. The only difference seen in this age group was a tendency for injection-site pain to persist somewhat longer in Menjugate recipients (72 hours) than in polysaccharide vaccine recipients (48 hours). This difference may be due to the aluminum hydroxide adjuvant, which is present in Menjugate but not the polysaccharide vaccine.

Infants:

Table 2 presents a summary of clinical safety data from two clinical studies in infants who received up to three immunizations with Menjugate, beginning at the age of two months.

Table 2- Summary of Local and Systemic Post immunization Reactions Within 7 Days Following 1, 2 or 3 Injections of Menjugate

	Percentage of Subjects	
	UK (Multicenter) (N=467)	Canada (Multicenter) (N=175)
Age at First Immunization	2 months	2 months
Schedule	3 doses 1 month apart	3 doses 2 months apart
Concomitant Vaccine	DTP, HIB, OPV	DTaP, HIB, IPV (PENTACEL)
Local Reactions:		
Tenderness	31%	22%
Erythema>25 mm	7%	0%
Induration>25 mm	4%	0%
Systemic Reactions:		
Irritability	81%	68%
Sleepiness	69%	54%
Change in Eating Habits	46%	39%
Diarrhea	43%	28%
Vomiting	34%	19%
Rash	16%	--*
Temp≥38°C	4%	21%
High-pitched crying	38%	--*
Persistent crying	16%	4%

* Data not collected

In a randomized, controlled clinical study performed in infants at three centers in Canada, the profile for Menjugate administered at 2, 4, and 6 months of age with concomitant PENTACEL (DTaP/Hib/IPV) was similar to that observed in earlier infant studies (See Tables 3 and 4 below). The frequency of two local adverse events, induration and erythema, was higher in Menjugate recipients than in the control HBV vaccine subjects, however the incidence of these reactions was lower among Menjugate or HBV subjects than following the routine vaccine (DTaP/Hib/IPV) in these same subjects. These differences between the Menjugate and HBV groups may in part be related to the lower dose of aluminum hydroxide in the HBV vaccine relative to the Menjugate vaccine (i.e., 0.5 mg per dose in HBV compared with 1 mg per dose in Menjugate). The most frequently reported systemic reactions were irritability, analgesic/antipyretic medication use, sleepiness and change in eating habits, which were reported with similar frequency in Menjugate and HBV vaccine subjects.

Table 3- Local Reactogenicity Within 7 Days Following Any Immunization Infant Study - Canada (Multicenter)

Local Reactions	Menjugate group N=175		HBV group N=176		P-value MenC vs HBV group	
	Menjugate	PENTACEL	HBV	PENTACEL	Study Vaccine	PENTACEL
Tenderness (Any) (Cried when injected leg moved)	38 (22%) 0	53 (30%) 3 (2%)	31 (18%) 0	35 (20%) 0	.33	.025
Erythema (Any) >25 mm	55 (31%) 0	67 (38%) 5 (3%)	33 (19%) 0	63 (36%) 4 (3%)	.006	.63
Induration (Any) >25 mm	42 (24%) 0	65 (37%) 6 (3%)	19 (11%) 1 (1%)	70 (40%) 2 (1%)	.001	.61

Table 4- Systemic Reactogenicity Within 7 Days Following Any Immunization Infant Study - Canada (Multicenter)

Systemic Reactions	Menjugate group N=175	HBV group N=176	P-value
Change in Eating Habits	68 (39%)	63 (36%)	.55
Sleepiness	94 (54%)	98 (56%)	.71
Persistent Crying	7 (4%)	4 (2%)	.35
Irritability	119 (68%)	124 (70%)	.62
Vomiting	34 (19%)	39 (22%)	.53
Diarrhea	49 (28%)	44 (25%)	.52
Rectal temp $\geq 38^{\circ}\text{C}$	37 (21%)	47 (27%)	.22
Analgesic/antipyretic medication required	96 (55%)	105 (60%)	.36

Less Common Clinical Trial Adverse Drug Reactions (<1%)

In clinical trials of Menjugate, approximately 6700 infants through adults were evaluated/monitored for the occurrence of serious adverse experiences (SAEs). There were four SAEs which were considered to be at least possibly related to vaccine. These were one report each of: hypotonia, screaming syndrome, maculopapular rash and agitation, all of which occurred in an open label infant study conducted in the United Kingdom (UK), in which Menjugate was administered concomitantly with DTP, Hib and OPV vaccines. Because these reactions have been reported previously in conjunction with DTP vaccines alone, a causal relationship between these experiences and Menjugate administration cannot be established.

Post-Market Adverse Drug Reactions

The most commonly reported suspected reactions in post marketing surveillance include dizziness, pyrexia, headache, nausea, vomiting and faints.

The frequencies given below are based on spontaneous reporting rates, for this and other Meningococcal C Conjugate vaccines and have been calculated using the number of reports received as the numerator and the total number of doses distributed as the denominator.

Immune System Disorders:

Very rare (<0.01%): lymphadenopathy, anaphylaxis including anaphylactic shock, hypersensitivity reactions including bronchospasm, facial edema and angioedema.

Neurologic:

Very rare (<0.01%): dizziness, convulsions including febrile convulsions, faints, hypoaesthesia and paresthesia, hypotonia.

There have been very rare reports of seizures following Menjugate vaccination; individuals have usually recovered rapidly. Some of the reported seizures may have been faints. The reporting rate of seizures was below the background rate of epilepsy in children. In infants seizures were usually associated with fever and were likely to be febrile convulsions.

There have been very rare reports of visual disturbances and photophobia following vaccination with Meningococcal group C conjugate vaccines, usually in conjunction with other neurological symptoms like headache and dizziness.

Skin and Subcutaneous tissue disorders:

Very rare (<0.01%): rash, urticaria, pruritus, purpura, erythema multiforme and Stevens-Johnson Syndrome.

Gastrointestinal:

Very rare (<0.01%): nausea, vomiting and diarrhea.

Musculoskeletal, connective tissue and bone disorders:

Very rare (<0.01%): myalgia and arthralgia.

Renal:

Relapse of nephrotic syndrome has been reported in association with Meningococcal group C conjugate vaccines.

DRUG INTERACTIONS

Drug-Drug Interactions

Administration of Menjugate at the same time as IPV, DTP, Hib, DTaP, DT, Td, MMR and HBV¹ vaccines or with OPV does not reduce immunological responses to any of these other antigens.² Menjugate should not however, be administered via the same injection as other vaccines. Minor variations in GMT antibody titers were observed between studies; however, the clinical significance, if any, of these observations is not established.

Table 5 presents data on the immunological response of infants to concomitant vaccines, as measured one month after the third dose of Menjugate or HBV vaccine.

Table 5- Response to Routine Infant Concomitant Vaccine (PENTACEL) Antigens Among Menjugate vs HBV Vaccine Recipients Measured at 1 Month After Third Dose Infant Study - Canada (Multicenter)³

Concomitant Vaccine Antigen, Measure of Response		Group		
		MenC	HBV	P-value
		N=64	N=61	
Polio type I	% with antibody titers \geq 1:8	97%	95%	.53
Polio type II	% with antibody titers \geq 1:8	98%	100%	.32
Polio type III	% with antibody titers \geq 1:8	98%	98%	.94
Diphtheria toxin antibody responses	GMT (IU/mL)	4.7	1.9	< .001
	% \geq 0.10 IU/mL	100%	100%	1.0
Tetanus antibody response	GMT (IU/mL)	2.4	2.5	.76
	% \geq 0.1 IU/mL	100%	100%	1.0
		N=91	N=89	
Anti-pertussis with 69K antigen	GMT (EU/mL)	31	36	.29
Anti-pertussis with FHA antigen	GMT (EU/mL)	26	31	.12
Anti-pertussis with PT antigen	GMT (EU/mL)	23	25	.31
		N=148	N=148	
PRP-T/Hib antibody responses	GMT (μ g/mL)	3.1	3.7	.28
	% \geq 1.0 μ g/mL	81%	83%	.40

DOSAGE AND ADMINISTRATION

Dosing Considerations

There are no data on the use of different meningococcal group C conjugate vaccines within the primary series or for boosting. Whenever possible, the same vaccine should be used throughout.

Recommended Dose and Dosage Adjustment

Primary Immunization: Infants 2-12 months should receive 3 doses of 0.5 mL each, with an interval of at least 1 month between doses.

Children older than 12 months, adolescents and adults should receive a single dose of 0.5 mL.

Booster doses

It is recommended that a booster dose should be given after completion of the primary

immunization series in infants. The timing of this booster should be in accordance with NACI recommendations. The need for booster doses in subjects primed with a single dose (i.e. aged 12 months or more when first immunized) has not yet been established.

For further information on responses to booster doses and on co-administration with other childhood vaccines please see DRUG INTERACTIONS and CLINICAL TRIALS.

Administration

Do not inject intravenously, subcutaneously or intradermally. Menjugate is to be administered by deep intramuscular injection only, preferably in the anterolateral thigh in infants and in the deltoid region in older children, adolescents and adults.

Care must be taken to ensure the vaccine is not injected into a blood vessel.

Menjugate should not be mixed with other vaccines in the same syringe. Separate injection sites should be used if more than one vaccine is being administered on the same date.

Syringe:

Gently shake the syringe containing the vaccine before administration. Remove the syringe tip cap and fit a suitable needle. The vaccine should be visually inspected for particulate matter and discoloration prior to administration. Ensure that no air bubbles are present in the syringe before injecting the vaccine. In the event of any foreign particulate matter and/or variation of physical aspect being observed, discard the vaccine.

Vial:

Gently shake the vaccine vial. Using a syringe and a suitable needle (21G, 1 ½ inch (40 mm) length) withdraw the entire content of vial. Prior to injection, change the needle for one suitable for the administration. The vaccine should be visually inspected for particulate matter and discoloration prior to administration. Ensure that no air bubbles are present in the syringe before injecting the vaccine. In the event of any foreign particulate matter and/or variation of physical aspect being observed, discard the vaccine.

OVERDOSAGE

There is no experience of overdosage with Menjugate.

ACTION AND CLINICAL PHARMACOLOGY

Menjugate is intended for the prevention of meningitis and/or septicemia caused by *Neisseria meningitidis* group C in infants and older age groups. Menjugate is composed of meningococcal group C oligosaccharides conjugated to a protein carrier, a non-toxic mutant of diphtheria toxin, CRM197. In the final vaccine, aluminum hydroxide is used as an adjuvant.

As shown in clinical trials, Menjugate is highly immunogenic^{4,5} and induces protective levels of bactericidal antibodies⁶⁻⁸ in a significant number of subjects after vaccination (See CLINICAL TRIALS).

Compared to licensed unconjugated polysaccharide vaccines, the primary immune response induced by Menjugate is superior in children (from 12 months of age) and adolescents,⁹ and is comparable in adults.

Additionally, unlike unconjugated polysaccharide vaccines,¹⁰ Menjugate has been shown to induce immunologic memory in infants and children from 12 months of age.^{2,11,12}

No pharmacodynamic or pharmacokinetic studies have been conducted with Menjugate, in accordance with its status as a vaccine. Several immunogenicity studies were conducted in animals, showing that Menjugate induced antibody titers that were dose dependent. (See Detailed Pharmacology).

STORAGE AND STABILITY

Primary storage condition: Store under refrigeration (+2°C to +8°C). Do not freeze. Protect from exposure to light. Stability studies have indicated that the product has a shelf life of 36 months.

DOSAGE FORMS, COMPOSITION AND PACKAGING

Dosage Forms

Menjugate Liquid is supplied as a 0.5 mL dose, consisting of one vial or syringe of vaccine. The vaccine is a white opalescent suspension.

Composition

Each 0.5 mL dose of the vaccine contains:

Active Ingredients	
<i>Neisseria meningitidis</i> group C (strain C11) oligosaccharide - Conjugated to	10 µg
<i>Corynebacterium diphtheriae</i> ¹⁵ CRM-197 protein	12.5 to 25.0 µg
Adjuvant	
Aluminium hydroxide	1 mg
Excipients	
Histidine	0.776 mg
Sodium chloride	4.5 mg
Water for Injections	to 0.5 mL

Menjugate contains no preservative.

Packaging

Syringe

Menjugate Liquid is presented in a Type I glass syringe, with bromobutyl rubber stopper and tip cap (styrene butadiene Type II rubber), filled with 0.6 mL of vaccine.

Vial

Menjugate Liquid is presented in a Type I glass vial, with bromobutyl rubber stopper, filled with 0.6 mL of vaccine.

Vial of vaccine: Pack sizes containing one, five, and ten doses are available.

Syringe of vaccine: Pack sizes containing one, five and ten doses are registered.

Not all pack sizes may be marketed.

PART II: SCIENTIFIC INFORMATION

PHARMACEUTICAL INFORMATION


Drug Substance

Proper name:	Meningococcal Group C Conjugate Vaccine
Chemical name:	Not applicable
Molecular formula and molecular mass:	Not applicable
Structural formula:	Not applicable
Physicochemical properties:	Not applicable

CLINICAL TRIALS

As shown in clinical trials, Menjugate is highly immunogenic^{4,5} and induces protective levels of bactericidal antibodies⁶⁻⁸ in a significant number of subjects after vaccination. (See Figure 1 below.)^{9,11,13} Data from trials in infants using a 2, 3, 4 month schedule demonstrate that >98% of infants developed serum bactericidal antibody titers of at least 1:8 one month after the second and third dose.² A booster dose in the second year of life induces an anamnestic response.^{11,12,14} Compared to licensed unconjugated polysaccharide vaccines, the primary immune response induced by Menjugate is superior in children from 12 months of age and adolescents,⁹ and is comparable in adults. Additionally, unlike unconjugated polysaccharide vaccines,¹⁰ Menjugate has been shown to induce immunologic memory in infants and children from 12 months of age.^{2,11,12}

Figure 1: Bactericidal Responses 1 Month following Menjugate or Meningococcal Polysaccharide (MenPS) Vaccine by Age*



*Infants did not receive MenPS vaccine

1/GMT (95% CI) = reciprocal geometric mean titer (95% confidence interval)

In a randomised clinical study involving 989 children aged 12 months to 2 years the immunogenicity of Menjugate liquid formulation was compared with that of Menjugate lyophilised formulation produced with active substance from two different manufacturing sites. For Menjugate liquid formulation, the geometric mean of post-vaccination to baseline titers ratio (GMR) was 4.69 (4.01-5.49); for Menjugate lyophilised formulation the GMRs were 5.6 (4.79-6.54) and 6.34 (5.4-7.45). The antibody response induced by both formulations of Menjugate was comparable. This was demonstrated by the two-sided 95% CI for the corresponding vaccine group GMT ratios falling within the predefined equivalence interval (0.5 - 2.0) 28 days post-vaccination. At the same time point, the proportion of subjects with hSBA \geq 1:8 were 60% (54-65) for the liquid formulation, 63% (57-69) and 70% (64-76) for the lyophilized formulation. These results were consistent with the pooled rate observed in children aged 12-23 months in previous studies (63%, CI 60-67) with Menjugate lyophilized formulation.

There are no data in children 2-12 months of age with liquid formulation.

No pharmacodynamic or pharmacokinetic studies have been conducted with Menjugate, in

accordance with its status as a vaccine. Several immunogenicity studies were conducted in animals, showing that Menjugate induced antibody titers that were dose dependent. (See DETAILED PHARMACOLOGY).

DETAILED PHARMACOLOGY

No pharmacodynamics studies and no pharmacokinetics studies have been conducted with Menjugate, in accordance with its status as a vaccine.

Animal Immunogenicity

Multiple immunogenicity studies were conducted in mice and guinea pigs to evaluate different subcutaneous dose levels of Meningococcal Group C–CRM197 conjugate (MenC conjugate) with and without the adjuvant aluminum hydroxide. Results showed that the induced antibody titers were dose-dependent and were higher when the MenC conjugate was combined with the adjuvant. The number of animals responding increased with dose.

An immunogenicity study in infant baboons was also conducted¹⁶. Five infant baboons (1.5 to 4 months of age) per group were immunized with three intramuscular injections (0.5 mL each) of MenC conjugate in combination with a Hib conjugated vaccine reconstituted either with aluminum hydroxide or phosphate buffered saline (PBS). Injections were given four weeks apart.

Titers of serum antibodies to MenC conjugate were measured by an enzyme-linked immunosorbent assay (ELISA). Complement-mediated bactericidal activity against *N meningitidis* group C was measured in a bactericidal assay.

The MenC conjugate was well tolerated; there was no evidence of specific local reactions at the injection sites when animals were inspected four weeks after the injection. After two injections, high antibody titers to meningococcus group C (bactericidal) were observed in animals vaccinated with MenC conjugate adjuvanted with aluminum hydroxide, while MenC conjugate in PBS (no adjuvant) showed low titers at all times. The bactericidal assay response paralleled the ELISA results.

TOXICOLOGY

The nonclinical safety of intramuscular injections of single and multiple doses of Meningococcal Group C–CRM197 Conjugate Vaccine (equivalent to Menjugate[®]) was evaluated in rabbits. In each study, the human dose was administered to rabbits. Five studies were performed under Good Laboratory Practices (GLPs)¹⁷:

1	Single dose intramuscular toxicity study	The vaccine was well tolerated. Slight, transient intramuscular injection site inflammation was seen, and was consistent with other alum-containing products.
---	--	---

2	Repeat-dose subacute toxicity study (three intramuscular injections)	Three injections of the vaccine were well tolerated (no effects on clinical signs including skin irritation, body temperature or weight, hematology or serum chemistry, ophthalmologic exams, or organ weights). Microscopic examination of muscle injection sites indicated mild inflammation. The mild inflammatory changes generally resolved within 2 weeks.
3	Repeat-dose subacute toxicity study (five intramuscular injections)	Five injections of the vaccine were well tolerated (no effects on clinical signs including skin irritation, body temperature or weight, hematology or serum chemistry, ophthalmologic exams, or organ weights). Slight to moderate inflammatory changes in muscle injection sites were seen and were generally resolved within 2 weeks.
4	Dose range-finding study of the effects on embryo/fetal development (eight intramuscular injections)	Three doses of vaccine were administered prior to conception and five doses were given during gestation. No clinically relevant maternal toxicity was seen and there were no vaccine-related effects on the external appearance of fetuses.
5	Study of the effects on embryo/fetal development (eight intramuscular injections)	Three doses of vaccine were administered prior to conception and five doses were given during gestation. No clinically relevant maternal toxicity was seen and there were no vaccine-related effects on fetuses based on evaluation of external, soft tissue, skeletal, and developmental parameters.

Overall, the vaccine was well tolerated and was associated with injection site findings consistent with the administration of any alum-adjuvanted vaccine.

REFERENCES

1. Schmitt HJ, Steul KS, Borkowski A, Ceddia F, Ypma E, Knuf M, Two versus three doses of a meningococcal C conjugate vaccine concomitantly administered with a hexavalent DTaP-IPV-HBV/Hib vaccine in healthy infants. *Vaccine* 2008; 26: 2242-2252
2. MacLennan JM, Shackley F, Heath PT, et al. Safety, immunogenicity, and induction of immunologic memory by a serogroup C meningococcal conjugate vaccine in infants: a randomized controlled trial. *JAMA*. 2000;283:2795-2801.
3. Data on File
4. Granoff DM, Maslanka SE, Carlone GM, et al. A modified enzyme-linked immunosorbent assay for measurement of antibody responses to meningococcal C polysaccharide that correlate with bactericidal responses. *Clin Diagn Lab Immunol*. 1998;5:479-485
5. Maslanka SE, Gheesling LL, Libutti DE, et al. Standardization and a multilaboratory comparison of *Neisseria meningitidis* serogroup A and C serum bactericidal assays. *Clin Diag Lab Immunol*. 1997;4:156-167.
6. Mandrell RE, Azmi FH, Granoff DM. Complement-mediated bactericidal activity of human antibodies to poly α 2 \rightarrow 8 N-acetylneuraminic acid, the capsular polysaccharide of *Neisseria meningitidis* serogroup B. *J Infect Dis*. 1995;172:1279-1289.
7. Goldschneider I, Gotschlich EC, Artenstein MS. Human immunity to the meningococcus. I. The role of humoral antibodies. *J Exp Med*. 1969;129:1307-1326
8. Santos G, Wacknov B, Borrow R, et al. Comparison of bactericidal assays used to measure immunological responses to vaccines developed against *Neisseria meningitidis* serogroup C. *ICAAC 2000*; 17-20 October 2000; Toronto, Canada (accepted for poster presentation).
9. Choo S, Zuckerman J, Goilav C, et al. Immunogenicity and reactogenicity of a group C meningococcal conjugate vaccine compared with a group A+C meningococcal polysaccharide vaccine in adolescents in a randomised observer-blind controlled trial. *Vaccine*. 2000;18:2686-2692..
10. De Wals P, De Serres G, Niyonsenga T. Effectiveness of a mass immunization campaign against serogroup C meningococcal disease in Quebec. *JAMA*. 2001;285:177-181.
11. MacDonald NE, Halperin SA, Law BJ, et al. Induction of immunologic memory by conjugated versus plain meningococcal polysaccharide vaccine in toddlers: a randomized controlled trial. *JAMA*. 1998;280:1685-1689.
12. Richmond P, Borrow R, Goldblatt D, et al. Ability of 3 different meningococcal C conjugate vaccines to induce immunologic memory after a single dose in UK toddlers. *J Infect Dis*. 2001;183:160-163.
13. MacDonald NE, Halperin SA, Law BJ, et al. Can meningococcal C conjugate vaccine overcome immune hyporesponsiveness induced by previous administration of multiple doses of plain polysaccharide vaccine? *JAMA*. 2000;283:1826-1827.
14. Trotter CL, Andrews NJ, Kaczmarski EB, Miller E, Ramsay ME, Effectiveness of meningococcal serogroup C conjugate vaccine 4 years after introduction. *The Lancet*.

- 2004;364:365-367
15. Rappuoli R. Isolation and characterization of *Corynebacterium diphtheriae* nontandem double lysogens hyperproducing CRM197. *Appl Env Microbiol.* 1983;46:560-564.
 16. Granoff DM, McHugh YE, Raff HV, et al. MF59 adjuvant enhances antibody responses of infant baboons immunized with *Haemophilus influenzae* type b and *Neisseria meningitidis* group C oligosaccharide-CRM197 conjugate vaccine. *Infect Immun.* 1997;65:1710-1715.
 17. Data on File

PART III: CONSUMER INFORMATION

Menjugate[®] Liquid Meningococcal Group C Conjugate Vaccine

This leaflet is part III of a three-part "Product Monograph" published when Menjugate was approved for sale in Canada and is designed specifically for Consumers. This leaflet is a summary and will not tell you everything about Menjugate. Contact your doctor or pharmacist if you have any questions about the drug.

ABOUT THIS VACCINE

What the vaccine is used for:

Menjugate is a vaccine that is used to prevent disease caused by the bacteria named *Neisseria meningitidis* serogroup C. *Neisseria meningitidis* serogroup C bacteria can cause serious and sometimes life-threatening infections such as meningitis and septicaemia (blood poisoning).

What it does:

The vaccine works by causing your body to make its own protection (antibodies) against these *Neisseria meningitidis* group C bacteria.

This vaccine can only protect against meningococcal serogroup C bacteria. It cannot protect against other groups (strains) of meningococcal bacteria or against other causes of meningitis and septicaemia (blood poisoning).

If at any time you or your child experiences neck pain, neck stiffness or a dislike of light (photophobia), drowsiness or confusion, red or purple bruise-like spots that do not fade under pressure you should contact your doctor or local Accident and Emergency Department immediately.

This vaccine cannot cause meningitis C (meningococcal C disease).

This vaccine contains a protein (called CRM197) from the bacteria that cause diphtheria. Menjugate does not protect against Diphtheria disease. This means that you/your child should receive other vaccines to protect against diphtheria when these are due or when advised by your doctor.

When it should not be used:

Menjugate should not be used if the person receiving the vaccine has ever had an allergic reaction to:

- any ingredient in the vaccine or
- any component of the container.

What the medicinal ingredient is:

Each 0.5 ml dose of vaccine contains the following amount of active substance: 10 micrograms of *Neisseria meningitidis* group C (strain C11) oligosaccharide chemically joined to 12.5 to 25.0 micrograms of *Corynebacterium diphtheriae* CRM197 protein.

What the important nonmedicinal ingredients are:

Aluminum hydroxide adjuvant

For a full listing of nonmedicinal ingredients see Part 1 of the product monograph.

What dosage forms it comes in:

Menjugate Liquid is formulated as a suspension for injection

Menjugate Liquid is available in two presentations:

- Vial of vaccine
- Syringe of vaccine

WARNINGS AND PRECAUTIONS

BEFORE you use Menjugate talk to your doctor or pharmacist if you (or your child):

- have an infectious illness (for example, high temperature, sore throat, cough, cold or flu)
- have haemophilia or any problem that may stop your blood from clotting properly
- been told that you have a weak immune system for any reason. For example, you have been told that you do not produce antibodies very efficiently, or you are taking medicines that reduce your immunity to infections (such as anti-cancer drugs or high doses of corticosteroids)
- have had your spleen removed or been told that your spleen does not work as it should
- are over 65 years old
- suffer from a kidney disease in which large amounts of protein appear in the urine (called nephrotic syndrome)

Pregnancy and breast-feeding

If you are pregnant, likely to become pregnant or are breast-feeding, you must tell your doctor before Menjugate is given. Your doctor or nurse may still advise you to have Menjugate if you are at high risk of infection with meningococcal group C bacteria.

Driving and using machines

You may feel dizzy or experience some other side effects after the injection. These could interfere with your driving or operating machinery. Do not drive or operate machinery until you know how Menjugate affects you.

Important information about some of the ingredients of

Menjugate

Latex-sensitive individuals – for syringe presentation:

Although no natural rubber latex is detected in the syringe tip cap, the safe use of Menjugate in latex-sensitive individuals has not been established.

Please tell your doctor if you (or your child) ever had an allergic reaction to latex.

INTERACTIONS WITH THIS MEDICATION

Menjugate may be given at the same time as other vaccinations but any other injected vaccines should be given into a different arm or leg from the site of the Menjugate injection.

PROPER USE OF THIS VACCINE

Usual dose:

Menjugate will be given to you/your child by a health professional. The vaccine is usually given into the muscle of the thigh in infants, and into the shoulder muscle for older children, adolescents and adults. Your healthcare professional will take care to ensure the vaccine is not given into a blood vessel and will make sure that it is injected into muscle and not into the skin.

For children over the age of 12 months, adolescents and adults: a single dose (0.5 ml) of the vaccine is recommended.

For infants 2 months up to 12 months of age: a complete vaccination schedule in infants consists of three doses of Menjugate. The first dose is given to infants from 2 months of age. A gap of at least one month should occur between each of doses.

In order to maintain protection, a booster dose must be given after the infant course has been completed. Your doctor will advise you when your child should receive this.

Overdose:

Since Menjugate will be given by either a doctor or nurse, and each injection is a single dose of 0.5 millilitres, it is very unlikely that you (or your child) will be given too much vaccine. If you have any concerns about the amount of vaccine you (or your child) have been given, speak to your doctor or nurse.

SIDE EFFECTS AND WHAT TO DO ABOUT THEM

Like all medicines, Menjugate may cause side effects in some persons. If any side effects worry you, or if you/your child have unusual symptoms, please contact your health professional.

If a serious allergic reaction occurs (usually in less than 1 in 10,000 people) tell your doctor straight away or go immediately/ take your child to the nearest Accident and Emergency department because urgent medical help may be needed.

The possible symptoms of serious allergic reactions can include:

- Swelling of the lips, mouth, throat (which may cause difficulty in swallowing)
- Difficulty breathing with wheezing or coughing
- Rash and swelling of the hands, feet and ankles
- Loss of consciousness and
- Very low blood pressure.

These very rare reactions can occur immediately or very soon after the injection and there is usually a rapid recovery after the right treatment has been given.

Other allergic reactions may start some days after the vaccine is

given.

These include:

- rashes, sometimes with itching, purple skin spots or blotches,
- blistering rashes that may also cause ulcers in the mouth and around the genital organs.

The most common side effects reported during clinical trials usually lasted only one to two days and were not usually severe.

The side effects were:

Very common (in more than 1 in 10 people)

- In all age groups: redness, swelling and tenderness/pain at the injection site but these did not usually require further medical attention. Redness or swelling of at least 3 cm and tenderness causing discomfort with movement were rarely observed for more than 48 hours.
- Infants: being sick (vomiting)
- Infants and children (aged 12 through 23 months): irritability, drowsiness, difficulty sleeping, loss of appetite and diarrhoea.
- Secondary school children: headache
- Older children and adults: feeling generally unwell, muscle and joint pains
- Adults: feeling sick (nauseous)

Common (between 1 in 10 and 1 in 100 people)

- In all age groups: Fever (but rarely severe).
- Infants and children aged 12 through 23 months: crying
- Children aged 12 through 23 months : being sick (vomiting).
- Primary school children: headache

Other side effects reported during routine vaccination programmes include:

Very rare (less than 1 in 10,000 people)

Different age groups:

- enlarged lymph glands
- dizziness
- faints
- numbness
- tingling sensation or pins and needles
- temporarily reduced muscle tone
- visual disturbances and sensitivity to light. These have usually occurred together with headache and dizziness.

Although fits have been reported very rarely after vaccination with Menjugate, it is thought that some of these reports in teenagers and adults may have been faints. In infants and young children, fits were usually associated with high fever. The majority of people affected have recovered rapidly.

There have been very rare reports of relapse of a kidney disorder called nephrotic syndrome following vaccination with this type of vaccine.

This is not a complete list of side effects. For any unexpected effects while taking Menjugate, contact your doctor or pharmacist.

HOW TO STORE IT

Store Menjugate in a refrigerator (+2°C to +8°C). Do not freeze. Keep the product in the outer carton in order to protect from light. Keep out of reach of children.

REPORTING SUSPECTED SIDE EFFECTS

To monitor vaccine safety, the Public Health Agency of Canada collects information on serious and unexpected adverse events following vaccination. If you suspect you have had a serious or unexpected event following receipt of a vaccine you may notify the Public Health Agency of Canada :

By toll-free telephone: 866-844-0018

By toll-free fax: 866-844-5931

By email: caefi@phac-aspc.gc.ca

By regular mail:

Vaccine Safety Section

Centre for Immunization & Respiratory Infections Diseases,

Public Health Agency of Canada

100 Eglantine Driveway

A/L 0602C, Building #6

Tunney's Pasture

Ottawa, Ontario K1A 0K9

NOTE: Should you require information related to the management of the side effect, please contact your health care provider before notifying the Public Health Agency of Canada. The Public Health Agency of Canada does not provide medical advice.

MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be found at:

www.gsk.ca or by contacting the sponsor:

GlaxoSmithKline Inc.

7333 Mississauga Road

Mississauga, Ontario

L5N 6L4

1-800-387-7374

Last revised: September 15, 2015

©2015 GlaxoSmithKline Inc., All Rights Reserved.

Menjugate® is a Registered Trademark of GlaxoSmithKline Biologicals SA.