

PRODUCT MONOGRAPH

**BEXSERO**

Multicomponent Meningococcal B Vaccine (recombinant, adsorbed)

Suspension for Injection

Active Immunizing Agent for the Prevention of Meningococcal Disease

ATC Code: J07AH09

GlaxoSmithKline Inc.  
7333 Mississauga Road  
Mississauga, Ontario  
L5N 6L4

**Date of Initial Approval:**

December 06, 2013

**Submission Control No:** 237046

**Date of Revision:**

June 12, 2020

*© 2020 GSK group of companies or its licensor  
Trademarks are owned by or licensed to the GSK group of companies.  
PREVNAR is a trademark of Pfizer Canada ULC.*

## Table of Contents

<b>PART I: HEALTH PROFESSIONAL INFORMATION.....</b>	<b>3</b>
SUMMARY PRODUCT INFORMATION .....	3
DESCRIPTION.....	3
INDICATIONS AND CLINICAL USE.....	3
CONTRAINDICATIONS .....	4
WARNINGS AND PRECAUTIONS.....	4
ADVERSE REACTIONS.....	6
DRUG INTERACTIONS.....	23
DOSAGE AND ADMINISTRATION.....	24
OVERDOSAGE .....	25
ACTION AND CLINICAL PHARMACOLOGY .....	25
STORAGE AND STABILITY.....	27
SPECIAL HANDLING INSTRUCTIONS .....	27
DOSAGE FORMS, COMPOSITION AND PACKAGING .....	28
<b>PART II: SCIENTIFIC INFORMATION .....</b>	<b>29</b>
PHARMACEUTICAL INFORMATION.....	29
CLINICAL TRIALS.....	29
TOXICOLOGY .....	46
REFERENCES .....	46
<b>PART III: CONSUMER INFORMATION.....</b>	<b>49</b>

## BEXSERO

Multicomponent Meningococcal B Vaccine (recombinant, adsorbed)  
Suspension for Injection

### PART I: HEALTH PROFESSIONAL INFORMATION

#### SUMMARY PRODUCT INFORMATION

Route of Administration	Dosage Form / Strength	Clinically Relevant Nonmedicinal Ingredients
Intramuscular injection	<p>Suspension for injection.</p> <p>White opalescent liquid suspension.</p> <p>Recombinant <i>Neisseria meningitidis</i> serogroup B NHBA fusion protein, 50 mcg <sup>1,2,3</sup>.</p> <p>Recombinant <i>Neisseria meningitidis</i> serogroup B NadA protein, 50 mcg <sup>1,2,3</sup>.</p> <p>Recombinant <i>Neisseria meningitidis</i> serogroup B fHbp fusion protein, 50 mcg <sup>1,2,3</sup>.</p> <p>Outer membrane vesicles (OMV) from <i>Neisseria meningitidis</i> serogroup B strain NZ98/254, 25 mcg measured as amount of total protein containing the PorA P1.4 <sup>2</sup>.</p> <p><sup>1</sup> Produced in <i>E. coli</i> by recombinant DNA technology. <sup>2</sup> Adsorbed on aluminium hydroxide (0.5 mg aluminium). <sup>3</sup> NHBA (Neisserial Heparin Binding Antigen), NadA (<i>Neisseria</i> adhesin A), fHbp (factor H binding protein).</p>	Aluminium hydroxide, histidine, sodium chloride, sucrose, water for injection. Residue*: kanamycin.

\*From the manufacturing process.

#### DESCRIPTION

BEXSERO is a liquid vaccine that contains three purified *Neisseria meningitidis* serogroup B protein antigens: NadA (*Neisseria* adhesin A) as a single protein, NHBA (Neisserial Heparin Binding Antigen) as a fusion protein, fHbp (factor H Binding Protein) as a fusion protein and PorA P1.4 as the main antigen of Outer Membrane Vesicles (OMV) derived from *N. meningitidis* serogroup B, strain NZ 98/254. These four *N. meningitidis* serogroup B antigens are adsorbed on aluminium hydroxide.

#### INDICATIONS AND CLINICAL USE

BEXSERO is indicated for active immunization of individuals from 2 months through 25 years old against invasive disease caused by *N. meningitidis* serogroup B strains.

As the expression of antigens included in the vaccine is epidemiologically variable in circulating group B strains, meningococci that express them at sufficient levels are predicted to be susceptible to killing by vaccine-elicited antibodies (see section ACTION AND CLINICAL PHARMACOLOGY).

## **CONTRAINDICATIONS**

BEXSERO should not be administered to individuals who are hypersensitive to this vaccine or to any ingredient in the formulation or components of the container closure.

For a complete listing, see the Dosage Forms, Composition and Packaging section of the Product Monograph.

## **WARNINGS AND PRECAUTIONS**

### **General**

As with any vaccine, vaccination with BEXSERO may not protect all vaccine recipients. BEXSERO is not expected to provide protection against all circulating meningococcal serogroup B strains.

The vaccine antigens present in BEXSERO are also expressed by meningococci belonging to serogroups other than serogroup B. However, protection against invasive meningococcal disease (IMD) caused by other serogroups has not been studied. Therefore, protection against IMD caused by other serogroups should not be assumed.

Do not inject intravascularly, subcutaneously or intradermally.

As with all injectable vaccines, appropriate medical treatment and supervision should always be readily available in case of an anaphylactic event following the administration of the vaccine.

Anxiety-related reactions, including vasovagal reactions (syncope), hyperventilation or stress-related reactions may occur in association with vaccination as a psychogenic response to the needle injection (see section ADVERSE REACTIONS). It is important that procedures are in place to avoid injury from fainting.

There are limited data on the use of BEXSERO in patients with chronic medical conditions.

As with all injectable pediatric vaccines, the potential risk of apnoea and the need for respiratory monitoring for 48-72 hours should be considered when administering the primary immunization series to very premature infants (born  $\leq$  28 weeks of gestation) and particularly for those with a previous history of respiratory immaturity. As the benefit of vaccination is high in this group of infants, vaccination should not be withheld or delayed.

The tip cap of the syringe may contain natural rubber latex. Although the risk for developing allergic reactions is very small, health professional should consider the benefit-risk prior to administering this vaccine to individuals with known history of hypersensitivity to latex.

Kanamycin is used in early manufacturing process and is removed during the later stages of manufacture. If present, kanamycin levels in the final vaccine are less than 0.01 micrograms per dose. The safe use of BEXSERO in kanamycin-sensitive individuals has not been established.

### **Febrile Illness**

As with many other vaccines, the physician should be aware that a temperature elevation may occur following vaccination of infants and children (less than 2 years of age). Prophylactic administration of acetaminophen at the time of, and closely after vaccination, can reduce the incidence and intensity of post-vaccination febrile reactions in infants and children (less than 2 years of age).

Administration of BEXSERO should be postponed in individuals suffering from an acute severe febrile illness. However, the presence of a minor infection, such a cold, should not be a reason to defer vaccination.

### **Hematologic**

This vaccine should not be given to individuals with thrombocytopenia, hemophilia or any coagulation disorder that would contraindicate intramuscular injection, unless the potential benefit clearly outweighs the risk of administration.

### **Immune**

Individuals with impaired immune responsiveness, whether due to the use of immunosuppressive therapy, a genetic disorder, or other causes, may have reduced antibody response to active immunisation. Immunogenicity data are available in individuals with complement deficiencies, and in individuals with splenic dysfunction or asplenia (see CLINICAL TRIALS).

Individuals receiving treatment that inhibits terminal complement activation (for example, eculizumab) remain at increased risk of invasive disease caused by *Neisseria meningitidis* group B even following vaccination with BEXSERO.

### **Sexual Function/Reproduction**

There are no data on fertility in humans. No effects on fertility were observed in female rabbits receiving BEXSERO pre-mating and during pregnancy.

### **Special Populations**

**Pregnant Women:** Insufficient clinical data on exposed pregnancies are available. The potential risk for pregnant humans is unknown. Nevertheless, vaccination should not be withheld when there is a clear risk of exposure to meningococcal infection.

#### *Preclinical data*

Based on reproductive toxicology data in rabbits, BEXSERO is not predicted to affect pregnancy and parturition, or to increase the risk of embryofetal abnormalities.

**Nursing Women:** No data are available. The benefit-risk ratio must be examined before making the decision to immunise during breast-feeding.

### *Preclinical data*

In a rabbit study, no effects on postnatal development were observed in nursing offspring of vaccinated maternal animals through day 29 of lactation.

**Pediatrics (< 2 months of age):** No data are available.

**Adults:** Limited safety and immunogenicity data are available in individuals above 25 years of age. The safety and efficacy of BEXSERO in individuals above 50 years have not been established. See ADVERSE REACTIONS and CLINICAL TRIALS.

**Geriatrics (> 65 years of age):** No data are available.

## **ADVERSE REACTIONS**

### **Adverse Drug Reaction Overview**

In clinical trials, the most frequent local and systemic adverse reactions after vaccination with BEXSERO were tenderness, erythema, induration, fever, irritability, unusual crying, sleepiness in infants and children (less than 2 years of age), and pain, erythema, induration, malaise, headache, myalgia in adolescents and adults. Higher rates of antipyretic use were also reported for infants vaccinated with BEXSERO and routine vaccines. When BEXSERO was given alone, the frequency of fever was similar to that associated with routine infant vaccines administered during clinical trials. Most solicited reactions were mild or moderate in severity and transient. No increase in the incidence or severity of the adverse reactions was seen with subsequent doses of the vaccination series.

### **Adverse Drug Reactions in Clinical Trials**

*Because clinical trials are conducted under very specific conditions the adverse reaction rates observed in trials may not reflect the rates observed in practice, and should not be compared with the rates in the clinical trials of another vaccine. Adverse drug reaction information from clinical trials is useful for identifying vaccine-related adverse events and for approximating rates.*

The characterization of the safety profile of BEXSERO is based on data from 13 studies, including 9 randomized controlled clinical trials with 7,526 participants (from 2 months of age) who received BEXSERO. Among BEXSERO recipients, 4,843 were infants and children (less than 2 years of age), 1,503 were adolescents (11 to 17 years of age) and 1,180 were adults (above 17 years of age), respectively. Of the subjects who received the primary infant series of BEXSERO, 1,630 received an additional dose in the second year of life. Data for 988 infants and children (less than 2 years of age) and 594 children (2 to 10 years of age) exposed to BEXSERO in subsequent studies have additionally been evaluated.

Data on solicited local (tenderness/pain, erythema, swelling and induration) and systemic adverse reactions (change in eating habits, sleepiness, irritability, unusual crying, vomiting, diarrhea, rash, fever  $\geq 38^{\circ}\text{C}$  in infants and children (less than 2 years of age); myalgia, arthralgia, nausea, malaise, headache and fever in adolescents and adults) were collected in clinical studies on the day of vaccination and for the following 6 days after vaccination (days 1-7 after

vaccination). Most reactions were of a mild to moderate nature and resolved within 48 hours after vaccination with BEXSERO.

### **Infants and Children (less than 2 years of age)**

In clinical studies in infants, when BEXSERO was given alone, the frequency of fever was comparable to that associated with concomitant use of routine infant vaccines [Pneumococcal 7-valent Conjugate Vaccine, Diphtheria CRM197 Protein (PREVNAR) and diphtheria, tetanus, acellular pertussis, hepatitis B recombinant (adsorbed), inactivated poliomyelitis and adsorbed conjugated *Haemophilus influenzae* type b vaccine (INFANRIX HEXA)]. Fever occurred more frequently when BEXSERO was co-administered with routine infant vaccines. Higher rates of antipyretic use were also reported for infants vaccinated with BEXSERO and routine vaccines. When fever occurred, it generally followed a predictable pattern, with the majority resolving within 48 hours after vaccination.

The characterization of the safety profile of BEXSERO in the infant and children (less than 2 years of age) populations was based primarily on data from 3 studies: V72P12 and V72P13 in infants 2 months of age, and V72P13E1 in children 12 to 13 months of age. In studies V72P12 and V72P13, the main schedule investigated was a three-dose primary series of BEXSERO administered at 2, 4 and 6 months of age. A three-dose accelerated schedule given at 2, 3 and 4 months of age was also evaluated in V72P12. BEXSERO was routinely administered with infant vaccines, INFANRIX HEXA and PREVNAR, except for one group of subjects in study V72P12 who received BEXSERO alone at 2, 4 and 6 months of age and the routine vaccines at 3, 5 and 7 months of age. In study V72P13E1, which was an extension of V72P13, subjects who previously received BEXSERO at the 2, 4, 6-month schedule received a fourth dose of BEXSERO at 12 months of age; control subjects who received only the routine infant vaccines in V72P13 (vaccine naive) were vaccinated with a two-dose catch-up schedule of BEXSERO at either 12 and 14 or 13 and 15 months of age.

### **Solicited Adverse Reactions**

#### ***3+1 Infant Schedule***

Data on local and systemic reactions after vaccination of infants with BEXSERO at 2, 4 and 6 months of age are shown in Table 1 and Table 2. Most of the reactions were transient and there was no clear trend of increasing frequency with subsequent doses. The reactogenicity profile was comparable for BEXSERO administered at the 2, 3, 4-month schedule.

Fever ( $\geq 38$  °C) was more frequently reported following vaccination with BEXSERO concomitantly with routine vaccines, compared with meningococcal C conjugate vaccine (MENJUGATE) with concomitant routine vaccines, or routine vaccinations only (Table 2). The onset of fever in the majority of BEXSERO recipients occurred within 6 hours of vaccination and the duration of the fever was transient, resolving within 48 hours after vaccination. This pattern was consistent for all three BEXSERO doses. There was a trend for subjects to have a higher probability of developing fever at a subsequent dose of BEXSERO if the subject experienced fever at the preceding dose(s).

More subjects used antipyretics after vaccination with BEXSERO and routine vaccines simultaneously than did those vaccinated with either BEXSERO or routine vaccines alone (Table

2). Even though fever rates were higher in subjects vaccinated with BEXSERO and concomitant vaccines, rates for fever in which a medical visit was sought were low and comparable to recipients of MENJUGATE with routine vaccines and routine vaccines only (Table 2).

Systemic reaction rates were comparable between the 2, 4, 6-month and 2, 3, 4-month schedules for recipients of BEXSERO with routine vaccines. For those subjects who received BEXSERO alone in the 2, 4, 6-month schedule without concomitant vaccines, fever rates were reduced (26% to 41% across the three doses) and comparable to the rates in subjects receiving only the routine infant vaccines.

**Table 1 - Percentage of Infants Experiencing Local Reactions on Days 1-7 Following Vaccination with BEXSERO and Routine Vaccines (INFANRIX HEXA, PREVNAR) at 2, 4, and 6 Months of Age**

		Percentage of Subjects With Injection Site Reactions (Severe or >100mm <sup>a</sup> )		
	Dose	BEXSERO Site <sup>b</sup>	INFANRIX HEXA Site <sup>c</sup>	Prevnar Site <sup>d</sup>
	1	N=3,101	N=3,102	N=3,102
	2	N=3,044	N=3,047	N=3,047
	3	N=3,019	N=3,023	N=3,022
Tenderness	1	66(14)	56(11)	54(11)
	2	66(14)	57(11)	55(11)
	3	65(14)	58(12)	56(11)
Erythema	1	60(<1)	46(0)	41(0)
	2	63(0)	57(0)	49(0)
	3	64(<1)	58(0)	52(0)
Induration	1	51(0)	33(0)	25(0)
	2	54(0)	47(0)	35(0)
	3	55(0)	49(0)	36(0)
Swelling	1	26(<1)	16(0)	13(0)
	2	27(<1)	21(0)	17(0)
	3	31(<1)	23(0)	19(0)

<sup>a</sup> Severe tenderness - cried when injected limb was moved; erythema, induration and swelling - >100 mm;

<sup>b</sup> BEXSERO: combined data of BEXSERO (studies V72P12 and V72P13) administered concomitantly with routine vaccines (INFANRIX HEXA, PREVNAR) in a 2, 4, 6-month schedule;

<sup>c</sup> INFANRIX HEXA vaccine administered in a 2, 4, 6-month schedule (studies V72P12 and V72P13);

<sup>d</sup> PREVNAR vaccine administered in a 2, 4, 6-month schedule (studies V72P12 and V72P13).



**Table 2 - Percentage of Infants Experiencing Systemic Reactions on Days 1-7 Following Vaccination with BEXSERO and Routine Vaccines (INFANRIX HEXA, PREVNAR) at 2, 4 and 6 Months of Age**

	Percentage of Subjects With Systemic (Severe <sup>a</sup> ) Reactions			
	Dose	BEXSERO+Routine Vaccines Group <sup>b</sup>	MENJUGATE+Routine Vaccines Group <sup>c</sup>	Routine Vaccines Only Group <sup>d</sup>
	1	N=3,102	N=490	N=659
	2	N=3,046-3,048	N=478-479	N=654
	3	N=3,023-3,024	N=470-471	N=651
Change Eat. Habits	1	51(3)	31(1)	30(2)
	2	44(3)	32(1)	25(<1)
	3	43(3)	29(1)	25(2)
Sleepiness	1	72(3)	58(4)	56(2)
	2	64(2)	45(1)	42(<1)
	3	53(1)	35(1)	32(<1)
Vomiting	1	13(1)	11(<1)	7(<1)
	2	13(<1)	11(1)	6(<1)
	3	12(<1)	9(<1)	7(<1)
Diarrhea	1	24(1)	20(1)	17(1)
	2	22(1)	15(<1)	17(<1)
	3	18(1)	13(1)	12(<1)
Irritability	1	79(6)	55(3)	61(2)
	2	79(7)	58(4)	62(3)
	3	76(6)	49(3)	54(1)
Unusual Crying	1	69(5)	52(3)	41(2)
	2	66(5)	50(4)	40(2)
	3	56(4)	39(3)	30(2)
Rash (Urticarial)	1	5(1)	4(<1)	3(1)
	2	6(2)	4(<1)	5(1)
	3	5(1)	3(0)	5(1)
<b>Other Solicited Outcomes</b>				
Fever $\geq 38^{\circ}\text{C}^{\text{e}}$ ( $\geq 40^{\circ}\text{C}$ )	1	75(<1)	46 (0)	44(<1)
	2	79(1)	63(<1)	59(<1)
	3	69(1)	42(0)	50(1)
Analgesic/ Antipyretic Medication use <sup>f</sup>	1	75	40	43
	2	81	52	52
	3	71	36	45
Medically Attended Fever <sup>g</sup>	1	1	1	1
	2	1	1	<1
	3	1	2	1

<sup>a</sup> Definition of severe: change in eating habits-missed >2 feeds; sleepiness-sleeps most of the time, hard to arouse; vomiting-little/no intake for more prolonged time; diarrhea -  $\geq 6$  liquid stools, no solid consistency; Irritability-unable to console; Unusual crying-unusual, high pitched, screaming, unlike the child's normal crying, that persists for  $\geq 3$  hours;

<sup>b</sup> BEXSERO+Routine Vaccines Group: combined data (studies V72P12 and V72P13) from BEXSERO administered concomitantly with routine vaccines (INFANRIX HEXA, PREVNAR) at a 2, 4, 6-month schedule;

<sup>c</sup> MENJUGATE+Routine Vaccines Group: data from MENJUGATE administered concomitantly with routine vaccines (INFANRIX HEXA, PREVNAR) from study V72P13 at a 2, 4, 6-month schedule;

<sup>d</sup> Routine Vaccines Only Group: data from routine vaccines (INFANRIX HEXA, PREVNAR) administered at a 2, 4, 6-month schedule from study V72P13;

<sup>e</sup> Fever is based on actual temperature recorded with no adjustment for route of measurement.

Body temperature was measured mainly by the rectal route in study V72P13; in study V72P12 body temperature was measured by both the rectal and axillary routes (30-31% rectal, 58-61% axillary);

<sup>f</sup> Percentage of subjects who were treated with analgesic or antipyretic medication during the day 1-7 time period after study vaccination;

<sup>g</sup> Percentage of subjects who had fever for which a medical visit was sought during the day 1-7 time period after study vaccination.

In an additional study, V72P16, BEXSERO was administered with INFANRIX HEXA and PREVNAR at 2, 3 and 4 months of age, with or without prophylactic acetaminophen. Data from this study showed that there is a statistically significant reduction in the percentage of subjects reporting fever both within 3 days and 7 days after vaccination when prophylactic acetaminophen treatment is adopted, without impacting the immune responses (see Part II, Immunogenicity Data).

Data on local and systemic reactions in children less than 2 years of age receiving either a fourth dose (booster) or two catch-up doses of BEXSERO are shown in Table 3 and Table 4. Additional data for a fourth dose of BEXSERO at 12 months of age in study V72P16 (after three doses at 2, 3 and 4 months of age) and at 12, 18 or 24 months of age in study V72P12E1 (after three doses at either 2, 4 and 6 months of age or 2, 3 and 4 months of age) confirmed these results. Data for a two-dose catch-up schedule of BEXSERO at either 12 and 14 or 18 and 20 months of age in control subjects who received only the routine infant vaccines in V72P12 are also in line with these observations.

In general, the majority of the local and systemic reactions following either a fourth dose or two-dose catch-up series of BEXSERO were transient, and most were mild or moderate in severity. Reactions (except tenderness) did not become more frequent after the second catch-up dose of BEXSERO.

**Table 3 - Percentage of Children (less than 2 years of age) Experiencing Local Reactions on Days 1-7 Following Vaccination with a Fourth Dose of BEXSERO at 12 Months of Age or with Two Catch-Up Doses of BEXSERO at 13 and 15 or 12 and 14 Months of Age, With or Without Concomitant PRIORIX-TETRA**

Percentage of Subjects With Injection Site Reactions (Severe or >50mm <sup>a</sup> )					
		4 <sup>th</sup> Dose of BEXSERO		Two Catch-up Doses of BEXSERO	
Schedule	Dose	BEXSERO with PRIORIX-TETRA at 12 mos.	BEXSERO at 12mos.	Dose 1: PRIORIX-TETRA at 12 mos.	Dose 1: BEXSERO with PRIORIX-TETRA at 12 mos.
		N=765	N=789	Dose 2: BEXSERO at 13 mos.	Dose 2: BEXSERO at 14 mos.
				Dose 3: BEXSERO at 15 mos.	N=117
				N=281	
Tenderness	1	71(14)	71(15)	20(1) <sup>b</sup>	57(10)
	2	-	-	56(10)	67(18)
	3	-	-	66(16)	-
Erythema	1	66(8)	68(7)	42(0) <sup>b</sup>	68(2)
	2	-	-	62(1)	60(2)
	3	-	-	58(3)	-
Induration	1	51(4)	54(3)	19(0) <sup>b</sup>	49(1)
	2	-	-	40(<1)	46(<1)
	3	-	-	42(<1)	-
Swelling	1	37(6)	36(5)	9(0) <sup>b</sup>	31(1)
	2	-	-	29(1)	28(1)
	3	-	-	30(3)	-

<sup>a</sup> Severe tenderness-cried when injected limb was moved; erythema, induration and swelling - >50 mm;

<sup>b</sup> Local reactions at the PRIORIX-TETRA injection site; mos: months.

**Table 4 - Percentage of Children (less than 2 years of age) Experiencing Systemic Reactions on Days 1-7 Following Vaccination with a Fourth Dose of BEXSERO at 12 Months of Age or with Two Catch-Up Doses of BEXSERO at 13 and 15 or 12 and 14 Months of Age, With or Without Concomitant PRIORIX-TETRA**

		Percentage of Subjects With Systemic Reactions (Severe <sup>a</sup> )			
		4 <sup>th</sup> Dose of BEXSERO		Two Catch-up Doses of BEXSERO	
Schedule	Dose	BEXSERO	BEXSERO	Dose 1:	Dose 1:
		with PRIORIX-TETRA at 12 mos.	at 12 mos.	PRIORIX-TETRA at 12 mos.	BEXSERO with PRIORIX-TETRA at 12 mos.
				Dose 2:	Dose 2:
				BEXSERO at 13 mos.	BEXSERO at 14 mos.
				Dose 3:	
				BEXSERO at 15 mos.	
		N=764-765	N=789	N=274-284	N=116-117
Change in Eating Habits	1	41(2)	40(2)	25(1)	38(0)
	2	-	-	34(1)	37(3)
	3	-	-	30(2)	-
Sleepiness	1	47(1)	45(1)	30(<1)	47(1)
	2	-	-	39(1)	41(0)
	3	-	-	39(1)	-
Vomiting	1	7(<1)	5(<1)	7(0)	2(0)
	2	-	-	5(<1)	3(1)
	3	-	-	3(0)	-
Diarrhea	1	25(1)	20(1)	16(1)	29(0)
	2	-	-	15(0)	22(0)
	3	-	-	15(0)	-
Irritability	1	73(4)	68(3)	43(1)	70(3)
	2	-	-	60(2)	63(3)
	3	-	-	56(3)	-
Unusual Crying	1	43(2)	37(2)	19(1)	35(2)
	2	-	-	28(1)	36(3)
	3	-	-	27(1)	-
Rash (Urticarial)	1	7(3)	7(2)	7(3)	8(1)
	2	-	-	5(2)	3(2)
	3	-	-	4(1)	-
Fever $\geq 38^{\circ}\text{C}$ ( $\geq 40^{\circ}\text{C}$ )	1	47(1)	41(<1)	24(<1)	46(0)
	2	-	-	37(0)	43(0)
	3	-	-	35 (<1)	-
Antipyretic Medication use <sup>b</sup>	1	57	51	23	57
	2	-	-	42	50
	3	-	-	39	-
Med. Attended Fever <sup>c</sup>	1	1	2	1	1
	2	-	-	0	2
	3	-	-	1	-

<sup>a</sup> Definition of severe: change in eating habits-missed >2 feeds; sleepiness-sleeps most of the time, hard to arouse; vomiting-little/no intake for more prolonged time; diarrhea -  $\geq 6$  liquid stools, no solid consistency; Irritability-

unable to console; Unusual crying-unusual, high pitched, screaming, unlike the child's normal crying, that persists for  $\geq 3$  hours;

<sup>b</sup> Percentage of subjects who were treated with any antipyretic medication during the day 1-7 time period after study vaccination;

<sup>c</sup> Percentage of subjects who had fever for which a medical visit was sought during the day 1-7 time period after study vaccination;

mos: months

### ***2+1 Infant Schedule***

In an additional study, V72\_28, the occurrence of solicited local and systemic reactions in infants vaccinated with the 3+1-dose schedule (Group I received 3 primary doses of BEXSERO at 2½, 3½ and 5 months, followed by a booster at 11 months of age) were similar to the 2+1-dose schedule (Group II received 2 primary doses of BEXSERO at 3½ and 5 months, followed by a booster at 11 months of age). There were no new significant safety signals in the 2+1 BEXSERO vaccination group (Group II), apart from that of the known safety profile from the 3+1 dose schedule.

### ***Unsolicited Adverse Events***

#### ***3+1 Infant Schedule***

Between study day 1 and 7 months of age (1 month after the third dose), the percent of subjects experiencing unsolicited AEs in the BEXSERO with concomitant routine vaccines, MENJUGATE with concomitant routine vaccines, and routine vaccines only groups are shown in Table 5.

Overall, between study day 1 and 7 months of age, the most commonly reported AEs after any vaccination with BEXSERO were injection site reactions (most considered as possibly related to vaccination as these local reactions of induration, erythema, and swelling were solicited AEs continuing after the 7-day vaccination window) and upper respiratory tract infections (10%; mostly considered unrelated to vaccination).

**Table 5 - Overview of Unsolicited Adverse Events of BEXSERO Administered with Concomitant Routine Vaccines at 2, 4 and 6 Months of Age, Collected From Study Day 1 to 7 Months of Age, by Vaccine Group**

	Percentage of Subjects with Adverse Events		
	BEXSERO+Routine Vaccines Group <sup>a</sup> N=3,155	MENJUGATE +Routine Vaccines Group <sup>b</sup> N=488	Routine Vaccines Only Group <sup>c</sup> N=658
Any AEs	77	63	71
At least possibly related AEs	52	42	34
Serious AEs	4	3	3

<sup>a</sup> BEXSERO+Routine Vaccines Group: combined data (studies V72P6, V72P12 and V72P13) from BEXSERO administered concomitantly with routine vaccines (INFANRIX HEXA, PREVNAR) at a 2, 4, 6-month schedule;

<sup>b</sup> MENJUGATE+Routine Vaccines Group: data from MENJUGATE administered concomitantly with routine vaccines (INFANRIX HEXA, PREVNAR) at a 2, 4, 6-month schedule from study V72P13;

<sup>c</sup> Routine Vaccines Only Group: data from routine vaccines (INFANRIX HEXA, PREVNAR) administered at a 2, 4, 6-month schedule from study V72P13;

AEs: Adverse Events.

The percentage of subjects who experienced unsolicited AEs after a two-dose catch-up schedule of BEXSERO in vaccine naive children (in their second year of life) was 17% after the first dose and 15% after the second dose of the vaccine; 3% had AEs considered by the investigator to be at least possibly related to vaccination and <1% to 6% were considered serious. The most commonly reported AEs were local injection site reactions and systemic reactions that were originally solicited, but continued past day 7 after vaccination. All of the injection site reactions were at least possibly related to study vaccination. The percentage of subjects who experienced unsolicited AEs after the fourth dose of BEXSERO in the second year of life was 44% and 74% for subjects who received BEXSERO alone and those who received BEXSERO with concomitant PRIORIX-TETRA vaccine, respectively. The most commonly reported AE was injection site induration. Most of the other AEs were due to local injection site reactions and systemic reactions that were originally solicited, but continued past day 7 after the vaccination.

### ***2+1 Infant Schedule***

In an additional study, V72\_28, the percentage of subjects with unsolicited adverse events in infants vaccinated with the 3+1 dose schedule (Group I received 3 primary doses of BEXSERO at 2½, 3½ and 5 months, followed by a booster at 11 months of age) were similar to the 2+1 dose schedule (Group II received 2 primary doses of BEXSERO at 3½ and 5 months, followed by a booster at 11 months of age).

### ***Children (aged 2 years to 10 years)***

The characterization of the safety profile of BEXSERO in this population is based on data from 4 studies in more than 290 subjects: V72P12E1 and V72P13E2 in children 24 months of age, V72P6E1 and V72P9E1 in children 40 to 62 months of age. In all these studies, the schedule investigated was a two-dose primary series of BEXSERO administered with an interval of 2 months between doses.

Solicited Adverse Reactions

Data on local and systemic reactions following vaccination with BEXSERO in children 2 to 10 years of age are shown in Table 6 and Table 7. Most of the solicited reactions were mild or moderate in severity and transient. The percentages of subjects with fever ranged from 10% to 28% in this age group. These rates were lower with increasing age. Few children (0-3% of subjects) experienced body temperature  $\geq 40^{\circ}\text{C}$ . Fever associated with BEXSERO vaccination occurred early after vaccination, and was transient, with the majority resolving within 2 days. Medically attended fever events occurred in no more than 3% of children.

**Table 6 - Percentage of Children (2 to 10 Years of Age) Experiencing Local Reactions on Days 1-7 Following Vaccination with BEXSERO**

		Percentages of Subjects With Any (Severe) Reaction				
		24 to 26 months of age		40-44 months of age		60-62 months of age
Local Reaction	Study Dose	V72P12E1 (N=54)	V72P13E2 (N=112)	V72P6E1 (N=42)	V72P9E1 (N=41)	V72P9E1 (N=48)
	Pain	1	-	-	93 (21)	87 (8)
2*		-	-	85 (15)	95 (24)	91 (13)
Tenderness	1	87 (26)	88 (10)	-	-	-
	2*	81 (35)	89 (18)	-	-	-
Erythema	1	72 (2)	77 (0)	98 (0)	92 (0)	94 (0)
	2*	60 (0)	73 (1)	93 (0)	97 (0)	87 (0)
Induration	1	50 (0)	49 (0)	33 (0)	44 (0)	40 (0)
	2*	42 (0)	56 (0)	49 (0)	49 (0)	44 (0)
Swelling	1	35 (0)	31 (0)	48 (0)	26 (0)	46 (0)
	2*	37(0)	39 (0)	63 (0)	41 (0)	44 (0)

\* local reaction after second dose was evaluated in at least N=52 in study V72P12E1, N=108 in study V72P13E2, N=41 in study V72P6E1, N=37 in study V72P9E1 (40-44 months cohort) and N=45 in study V72P9E1 (60-62 months cohort).

**Table 7 - Percentage of Children (2 to 10 Years of Age) Experiencing Systemic Reactions Days 1-7 Following Vaccination with BEXSERO**

		Percentages of Subjects With Any (Severe) Reaction				
		24 to 26 months of age		40-44 months of age		60-62 months of age
Systemic Reaction	Study Dose	V72P12E1 (N=54)	V72P13E2 (N=112)	V72P6E1 (N=42)	V72P9E1 (N=39)	V72P9E1 (N=48)
	Change Eat. Habits	1	46 (2)	34 (0)	38 (2)	33 (3)
2*		40 (4)	36 (3)	34 (0)	35 (3)	22 (2)
Sleepiness	1	33 (2)	46 (0)	48 (5)	51 (8)	40 (6)
	2*	35 (0)	46 (3)	37 (2)	46 (8)	30 (0)
Vomiting	1	11 (2)	8 (0)	2 (0)	3 (0)	10 (0)
	2*	8 (2)	5 (0)	0	11 (0)	7 (0)
Diarrhea	1	37 (0)	13 (0)	14 (0)	5 (0)	4 (0)
	2*	13 (4)	12 (0)	2 (0)	5 (0)	4 (0)
Irritability	1	52 (7)	59 (2)	76 (7)	62 (0)	44 (4)
	2*	44 (4)	58 (5)	59 (5)	62 (5)	43 (2)
Unusual Crying	1	28 (2)	33 (1)	-	-	-
	2*	29 (4)	27 (3)	-	-	-
Headache	1	-	-	10 (0)	10 (0)	13 (2)
	2*	-	-	10 (2)	11 (0)	20 (0)
Arthralgia	1	-	-	31 (7)	23 (3)	31 (2)
	2*	-	-	22 (7)	19 (5)	33(2)
Rash	1	4 (0)	7 (3)	2 (2)	5 (0)	6 (0)
	2*	0	6 (0)	5 (0)	3 (0)	9 (2)
Fever [Body Temp. $\geq 38^{\circ}\text{C}$ ( $\geq 40^{\circ}\text{C}$ )]	1	28 (0)	21 (0)	10 (0)	15 (3)	10 (0)
	2*	25 (0)	26 (1)	12 (0)	11 (0)	11 (0)
Medical Attended Fever	1	2 (-)	0 (-)	0 (-)	0 (-)	0 (-)
	2*	0 (-)	2 (-)	0 (-)	3 (-)	0 (-)

\* systemic reaction after second dose was evaluated in at least N=52 in study V72P12E1, N=108 in study V72P13E2, N=41 in study V72P6E1, N=37 in study V72P9E1 (40-44 months cohort) and N=46 in study V72P9E1 (60-62 months cohort)



### Unsolicited Adverse Events

Table 8 provides an overview of unsolicited adverse events collected up to month 62 in children 2 to 10 years of age.

**Table 8 - Overview of Unsolicited Adverse Events in Children (2 to 10 Years of Age) After the Two-Dose Schedule of BEXSERO**

Study	Percentage of Subjects with Adverse Events				
	24 to 26 months of age		40-44 months of age		60-62 months of age
	V72P12E1	V72P13E2	V72P6E1	V72P9E1	V72P9E1
	N=55	N=112	N=42	N=41	N=48
Any AEs	75	86	43	61	38
Possibly related	29	36	14	15	27
SAEs	2	5	2	10	2

### Adolescents (aged 11 to 17 years)

The characterization of the safety profile of BEXSERO in the adolescent population aged 11 to 17 years was based on data from study V72P10. One, two or three doses of BEXSERO were administered to adolescents according to one of the following schedules: 0, 0-1, 0-2 or 0-1-2 months. The data supporting the safety and tolerability of the two-dose vaccination schedules for adolescents were generated from the first and second doses of the schedules investigated in this study.

### Solicited Adverse Reactions

Data on local and systemic reactions are shown in Table 9.

The frequency of reports for local and systemic reactions did not increase with the second dose of BEXSERO, and the majority of the reactions were transient.

Additional safety data on BEXSERO in adolescents relative to the administration of one dose at month 6 from study V72P10 and for 2 doses 1 month apart in study V72\_41 were in line with these observations.

**Table 9 - Percentage of Adolescents (aged 11-17 Years) Experiencing Local and Systemic Reactions on Days 1-7 Following Vaccination with BEXSERO**

	Percentage of Subjects With Any (Severe <sup>a</sup> ) Reaction		
	Study Schedule	V72P10 (Combined Month 0, 0-1, 0-2) 11-17 years of age	
	Dose	BEXSERO	Placebo <sup>b</sup>
	1	N=1,503	N=128
	2	N=1,039	N=124
<b>Local Reactions</b>			
Erythema	1	54(<1)	40(0)
	2	51(<1)	31(0)
Induration	1	40(<1)	27(0)
	2	40(<1)	23(0)
Swelling	1	39(<1)	20(0)
	2	38(1)	15(0)
Pain	1	91(17)	86(9)
	2	85(15)	71(9)
<b>Systemic Reactions</b>			
Malaise	1	56(7)	48(3)
	2	50(7)	35(2)
Myalgia	1	45(7)	41(4)
	2	40(6)	40(3)
Arthralgia	1	24(2)	19(0)
	2	21(3)	16(1)
Headache	1	46(5)	37(2)
	2	42(5)	33(3)
Nausea	1	19(1)	17(2)
	2	16(2)	15(1)
Fever $\geq 38^{\circ}\text{C}$ ( $\geq 40^{\circ}\text{C}$ )	1	3(0)	4(0)
	2	4(0)	2(0)
<b>Other Solicited Outcomes</b>			
Analgesic/antipyretic use <sup>c</sup>	1	35	20
	2	27	15
Stayed home due to reaction <sup>d</sup>	1	16	6
	2	11	3

<sup>a</sup> Severe erythema, induration and swelling - >100 mm; severe pain and systemic reactions - unable to perform normal daily activity;

<sup>b</sup> Placebo administered in month 0-1 schedule;

<sup>c</sup> Percentage of subjects who were treated with analgesic or antipyretic medication during the day 1-7 time period after study vaccination;

<sup>d</sup> Collected as yes or no

### Unsolicited Adverse Events

Table 10 provides an overview of unsolicited adverse events collected up to study Month 3 in adolescents who received BEXSERO in either the 0-1 month or 0-2 month schedule (study V72P10). The most commonly reported possibly or probably related unsolicited AEs were local injection site reactions (pain, induration, swelling) that continued past the day 7 observation period.

**Table 10 - Overview of Unsolicited Adverse Events Collected up to Month 3 in Adolescents (11 to 17 Years of Age), after the Two-Dose Schedule of BEXSERO**

	Percentage of Subjects with Adverse Events	
	0-1 Month Schedule <sup>a</sup>	0-2 Month Schedule <sup>b</sup>
	11 to 17 yoa N=748	11 to 17 yoa N=380
Any AEs	43	46
Possibly or probably related AEs	17	16
Serious AEs	1	1

<sup>a</sup> 0-1 schedule: BEXSERO was administered at months 0 and 1 in the 11 to 17 years of age (study V72P10);

<sup>b</sup> 0-2 schedule: BEXSERO was administered at months 0 and 2 in the 11 to 17 years of age (study V72P10);

yoa: years of age;

vs: versus;

AEs: Adverse Events.

### Adults

The characterization of the safety profile of BEXSERO in the adult population (above 17 years of age) was based on data from 7 studies, V72\_29, V72\_37, V72\_59, V72\_74, V72P4, V72P5, and V102\_03. Most of the subjects enrolled in the studies were between 18 and 24 years of age. In addition, there were 82 subjects 25 to 50 years of age and 3 subjects above 50 years of age. Two or three doses of BEXSERO were administered according to one of the following schedules: 0-1, 0-2, 0-1-2, or 0-2-6 months. The data supporting the safety and tolerability of the two-dose adult vaccination schedule were generated from data collected in the first and second doses, pooled across studies and stratified by age (18 to 24 years, and 25 years and older). Data on solicited local (pain, erythema, swelling and induration) and systemic (chills, nausea, malaise, myalgia, arthralgia, headache, fever, rash, fatigue, loss of appetite) reactions were collected in clinical studies on the day of vaccination and for the following 6 days after vaccination (days 1-7 after vaccination).

### Solicited Adverse Reactions

Percentage of adults experiencing local and systemic reactions on days 1 to 7 following each vaccination with BEXSERO are presented in Table 11 and Table 12. The incidence rate of the individual solicited local and systemic adverse events in subjects 18 to 24 years was generally higher in the BEXSERO group than in the control group. After both vaccinations, the most commonly reported local adverse reactions were pain, followed by erythema. The most commonly reported systemic adverse reactions were myalgia and headache. Most of the subjects reported solicited local and systemic adverse reactions which were mild to moderate in severity. The most commonly reported severe solicited reactions were pain and myalgia. Limited data from V72P4, V72P5, V72\_59 and V102\_03 in subjects above 25 years of age showed the same

trend in the most commonly reported solicited local and systemic adverse events and the severity of the adverse events as in subjects 18 to 24 years.

**Table 11 - Percentage of Adults (18 to 24 Years of Age) Experiencing Local Reactions on Days 1-7 Following Each Vaccination with BEXSERO, Pooled Across Studies, Two-Dose Schedules<sup>a</sup>**

Percentages of Subjects <sup>b</sup> With Any (Severe <sup>c</sup> ) Local Adverse Reactions					
Study	Studies with control arm V72_29 <sup>d</sup> + V102_03		Studies without control arm V72P4 + V72P5 + V72_59	Pooled across studies V72_29 <sup>d</sup> + V72_59 + V72P4 + V72P5 + V102_03	
	Dose	Control <sup>e</sup>	BEXSERO	BEXSERO	
	1	N = 402	N = 226	N = 12	N = 238
	2	N = 378	N = 216	N = 11	N = 227
Pain	1st	46 (0)	93 (10)	92 (17)	93 (11)
	2nd	49 (1)	88 (10)	91 (0)	88 (10)
Erythema	1st	25 (0)	41 (<1)	42 (0)	41 (<1)
	2nd	19 (0)	41 (1)	27 (0)	40 (1)
Induration	1st	11 (0)	27 (1)	42 (0)	28 (<1)
	2nd	9 (0)	23 (0)	36 (0)	24 (0)
Swelling <sup>b</sup>	1st	8 (0)	25 (1)	0 (0)	25 (1)
	2nd	7 (<1)	26 (0)	0 (0)	26 (0)

Notes: Only unsolicited adverse event data were collected in studies V72\_37 and V72\_74. Solicited safety set: all subjects in the exposed set who provided postvaccination reactogenicity data.

<sup>a</sup>0, 1 schedule: V72\_29, V72P5. All other studies had a 0, 2 schedule. For studies V72P4 and V72P5 an additional dose was given i.e., 3 doses were given compared to 2 doses in the other studies. The third dose was not included in the post-hoc analyses.

<sup>b</sup> Swelling was solicited for V72\_29 and V72\_59, not solicited in V102\_03, V72P4 or V72P5.

<sup>c</sup> Severe pain - unable to perform normal daily activity; severe erythema, induration and swelling - >100 mm.

<sup>d</sup> For study V72\_29 solicited local and systemic AEs were collected only in the immunogenicity subset, while all unsolicited AEs and SAEs were collected for all study participants.

<sup>e</sup> For study V72\_29, control group includes subjects that received IXIARO, as well as the Placebo/ACWY group where first vaccination was MenACWY, second vaccination was placebo. For study V102\_03, control group includes subjects whose first vaccination was placebo, second vaccination was MenACWY.

**Table 12 - Percentage of Adults (18 to 24 Years of Age) Experiencing Systemic Reactions on Days 1-7 Following Each Vaccination with BEXSERO, Pooled Across Studies, Two-Dose Schedules<sup>a</sup>**

Percentages of Subjects With Any (Severe <sup>b</sup> ) Systemic Adverse Reactions					
Study		Studies with control arm V72_29 <sup>c</sup> + V102_03		Studies without control arm V72P4 + V72P5 + V72_59	Pooled across studies V72_29 <sup>c</sup> + V72_59 + V72P4 + V72P5 + V102_03
	Dose	Control <sup>d</sup>	BEXSERO	BEXSERO	BEXSERO
	1	N = 402	N = 226	N = 12	N = 238
	2	N = 379	N = 215	N = 11	N = 226
Malaise <sup>e</sup>	1st	18 (1)	17 (1)	27 (0)	18 (<1)
	2nd	12 (1)	22 (2)	20 (0)	22 (2)
Nausea	1st	8 (<1)	13 (<1)	0 (0)	13 (<1)
	2nd	5 (1)	10 (<1)	9 (0)	10 (<1)
Myalgia	1st	45 (1)	70 (7)	33 (8)	68 (7)
	2nd	40 (1)	66 (7)	45 (0)	65 (6)
Arthralgia	1st	8 (<1)	12 (1)	25 (0)	13 (1)
	2nd	8 (1)	12 (1)	18 (0)	12 (1)
Headache	1st	27 (<1)	29 (1)	42 (0)	30 (1)
	2nd	17 (1)	22 (2)	27 (0)	22 (2)
Fever	1st	2 (0)	1 (0)	8 (0)	1 (0)
	2nd	1 (0)	2 (0)	0 (0)	2 (0)
Analgesic /antipyretic Use	1st	8 (-)	20 (-)	17 (-)	20 (-)
	2nd	6 (-)	23 (-)	0 (-)	21 (-)
Stayed home due to reaction	1st	2 (-)	3 (-)	27 (-)	4 (-)
	2nd	1(-)	6 (-)	20 (-)	7 (-)

Notes: Only unsolicited adverse event data were collected in studies V72\_37 and V72\_74. Solicited safety set: all subjects in the exposed set who provided postvaccination reactogenicity data.

<sup>a</sup> 0, 1 schedule: V72\_29, V72P5. All other studies had a 0, 2 schedule. For studies V72P4 and V72P5 an additional dose was given i.e., 3 doses were given compared to 2 doses in the other studies. The third dose was not included in the post-hoc analyses.

<sup>b</sup> Severe: malaise = unable to perform daily activity; nausea = leading to minimal to no oral intake; myalgia, arthralgia, headache = prevented daily activity; fever ≥40°C.

<sup>c</sup> For study V72\_29 solicited local and systemic AEs were collected only in the immunogenicity subset, while all unsolicited AEs and SAEs were collected for all study participants.

<sup>d</sup> For study V72\_29, control group includes subjects that received IXIARO, as well as the Placebo/ACWY group where first vaccination was MenACWY, second vaccination was placebo. For study V102\_03, control group includes subjects whose first vaccination was placebo, second vaccination was MenACWY.

<sup>e</sup> Malaise was solicited for V72\_29, V72P4 and V72P5 only.

### Unsolicited Adverse Events

Data on unsolicited AEs in adults (subjects 18 to 24 years) are shown in Table 13.

**Table 13 - Overview of Unsolicited Adverse Events Reported Up to 3 Months in Adults, Pooled Across Studies, Two-Dose Schedules<sup>a</sup>**

Study	Percentage of Subjects with Unsolicited Adverse Events
	V102_03 + V72_29 + V72P4 + V72P5 + V72_59 + V72_74
	<b>BEXSERO</b> N=1,048
Any AE	25
Possibly or probably related AEs	7
SAEs	2

Abbreviation: AE, adverse event, SAE, serious adverse event.

Notes: Unsolicited safety set: all subjects in the exposed set who provided post vaccination unsolicited AE records.

<sup>a</sup> 0, 1 schedule: V72\_29, V72P5. All other studies had a 0, 2 schedule. The reporting period here is up to 1 month following the second dose in a 0,2 dose schedule.

### **Less Common Drug Reactions Seen in Clinical Trials (<1%)**

Adverse reactions (following primary immunization or additional dose) considered as being at least possibly related to vaccination have been categorized by frequency.

Frequencies are defined as follows:

Uncommon:  $\geq 1/1,000$  to  $< 1/100$

Rare:  $\geq 1/10,000$  to  $< 1/1,000$

*Infants and Children (less than 2 years of age)*

General disorders and administration site conditions

Uncommon: fever ( $\geq 40^{\circ}\text{C}$ )

Nervous system disorders

Uncommon: seizures (including febrile seizures)

Skin and subcutaneous tissue disorders

Uncommon: eczema, urticaria

Vascular disorders

Uncommon: pallor (rare after booster)

Rare: Kawasaki syndrome

*Children (2 to 10 years of age)*

Nervous system disorders

Uncommon: seizures (including febrile seizures)

### Vascular disorders

Uncommon: pallor (rare after booster)

Rare: Kawasaki syndrome

### **Post-Market Adverse Drug Reactions**

In addition to reports in clinical trials, worldwide voluntary reports of adverse reactions received for BEXSERO since market introduction are listed below. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency.

### General disorders and administration site conditions

Fever (adolescents from 11 years of age and adults), injection site reactions (including extensive swelling of the vaccinated limb, blisters at or around the injection site and injection site nodule which may persist for more than one month)

### Immune system disorders

Allergic reactions (including anaphylactic reactions)

### Nervous system disorders

Hypotonic-hyporesponsive episode, syncope or vasovagal responses to injection

### Skin and subcutaneous tissue disorders

Rash (adolescents from 11 years of age and adults)

## **DRUG INTERACTIONS**

### **Drug-Drug Interactions**

BEXSERO can be given concomitantly with any of the following vaccine antigens, either as monovalent or as combination vaccines: diphtheria, tetanus, acellular pertussis, *Haemophilus influenzae* type b, inactivated poliomyelitis, hepatitis B, heptavalent pneumococcal conjugate, measles, mumps, rubella, varicella, and meningococcal groups A, C, W, Y conjugate. (see Part II, CLINICAL TRIALS, Immunogenicity Data, *Concomitant use of BEXSERO with routine vaccines*).

As higher percentages of subjects reported systemic reactions, including fever, change in eating habits, tenderness at the injection site and irritability, following BEXSERO given concomitantly with routine vaccines than after BEXSERO alone, separate vaccinations can be considered when possible. In addition, fever was mostly reported during the 1-4 days after vaccination with BEXSERO alone and during the 5-28 days after the MMRV vaccination alone.

Prophylactic use of acetaminophen reduces the incidence and severity of fever without affecting the immunogenicity of either BEXSERO or most antigens of routine vaccines. The effect of antipyretics other than acetaminophen on the immune response has not been studied.

Concomitant administration of BEXSERO with vaccines other than those mentioned above has not been studied.

When given concomitantly with other vaccines, BEXSERO should be administered at different injection site.

### **Drug-Lifestyle Interactions**

BEXSERO has no or negligible influence on the ability to drive and use machines. However, some of the effects mentioned under section ADVERSE REACTIONS may temporarily affect the ability to drive or use machines.

## **DOSAGE AND ADMINISTRATION**

### **Recommended Dose and Dosage Adjustment**

<b>Age at First Dose</b>	<b>Primary Doses</b>	<b>Intervals between Primary Doses</b>	<b>Booster</b>
Infants, 2 months to 5 months	<b>3+1 schedule:</b>		
	Three doses, each of 0.5 mL	Not less than 1 month	1 booster in 2 <sup>nd</sup> year of life with an interval of at least 6 months between the primary doses and booster.*
	<b>2+1 schedule:</b>		
	Two doses, each of 0.5 mL	Not less than 2 months	1 booster in 2 <sup>nd</sup> year of life with an interval of at least 6 months between the primary doses and booster.*
Infants, 6 months to 11 months	Two doses, each of 0.5 mL	Not less than 2 months	1 booster in 2 <sup>nd</sup> year of life with an interval of at least 2 months between the primary doses and booster.*
Children, 12 months to 23 months	Two doses, each of 0.5 mL	Not less than 2 months	1 booster given at an interval of 12 to 23 months between the primary series and booster.*
Children, Adolescents and Adults, 2 through 25 years	Two doses, each of 0.5 mL	Not less than 1 month	1 booster considered in individuals at continued risk of exposure to meningococcal disease, based on guideline recommendations.

\*The need for, and timing of further doses has not yet been determined (see Clinical Trials section).

Sufficient data are not available on the safety and effectiveness of using BEXSERO and other meningococcal group B vaccines interchangeably to complete the vaccination series. Therefore, it is recommended that subjects who receive a first dose of BEXSERO complete the vaccination course with BEXSERO.



### **Administration**

BEXSERO should be given by deep intramuscular injection, preferably in the anterolateral aspect of the thigh in infants or in the deltoid muscle region of the upper arm in older individuals.

Separate injection sites must be used if more than one vaccine is administered at the same time. The vaccine must not be injected intravenously, subcutaneously or intradermally and must not be mixed with other vaccines in the same syringe.

BEXSERO must not be mixed with other medicinal products.

### **OVERDOSAGE**

Experience of overdose is limited. In the event of overdose, monitoring of vital functions and possible symptomatic treatment is recommended.

For management of a suspected drug overdose, contact your regional Poison Control Centre
--

### **ACTION AND CLINICAL PHARMACOLOGY**

#### **Epidemiology Data**

*Neisseria meningitidis* is the bacterium which causes invasive meningococcal disease (IMD), an important cause of meningitis and septicaemia. Strains of *N. meningitidis* can be classified into 12 distinct serogroups based on the composition of their distinct polysaccharide capsules. Of the known serogroups, only 4 (B, C, W-135 and Y) are responsible for the vast majority of IMD within Canada. Following the implementation of meningococcal conjugate C immunization programs in Canada, and the corresponding decline in serogroup C IMD, serogroup B has become the leading cause of IMD.

From 2007-2011, the average annual number of IMD cases in Canada was close to 200 (range: 154 to 229). In the general population, approximately 60% of IMD cases have been attributed to serogroup B, with the greatest disease burden seen in young infants (<1 years) and children 1-4 years of age. A secondary peak of incidence occurs in adolescents and young adults 15-19 years of age. Pharyngeal carriage rates of meningococci are highest in the adolescent population and range from 20-30%. The case-fatality rates (CFRs) due to IMD vary by serogroup and age, and approximate 9% in Canada. Those who survive can face significant life-long sequelae. Sequelae were reported in 19% of serogroup B cases (7).

The potential of BEXSERO to protect against diverse invasive serogroup B strains isolated in Canada was studied using the *in-vitro* Meningococcal Antigen Typing System (MATS) that was specifically developed to provide an estimate of strain coverage by measuring the level of antigen expression on circulating strains together with a measure of antigenic diversity compared with the BEXSERO vaccine antigens. The MATS was established to predict the effect that the vaccine will have in market use by relating antigen profiles of different strains of meningococcal group B bacteria to killing of the strains in the serum bactericidal assay with human complement (hSBA). As the antigens, including NHBA, NadA, fHbp, and PorA P1.4, are variably expressed by different strains, meningococci that express at least one antigen at sufficient levels are susceptible to killing by vaccine-elicited antibodies (3). However, the MATS assay may provide variable estimates depending on the strains in circulation at a given time and geography. As

MATS coverage predictions are based on killing of meningococci by immune serum pools and not individual subject sera, this prediction is subject to certain limitations and its accuracy may only be verified upon vaccine use.

A survey of approximately 157 invasive serogroup B isolates collected during 2006-2009 by the IMPACT surveillance network (Immunization Monitoring Program ACTIVE) revealed that 66% of isolates had an appropriate type and sufficient antigen content and were predicted to be covered by BEXSERO with empirical variability in coverage from 43% to 78% (4).

Coverage for the Canadian hyper-endemic strains, the two most prevalent strains (sequence type ST-269 and ST-154) was 95% and 100%, with empirical variability in coverage ranging from 43% to 97% and 100% to 100%, respectively (4).

The hyper-endemic strain ST-269 emerged in the province of Quebec in 2003 and attributed to a steadily increasing incidence of serogroup B IMD with the Saguenay-Lac-Saint-Jean (SLSJ) region having the greatest serogroup B IMD incidence rates in Quebec. A short-term mass immunization campaign was initiated in May 2014 with BEXSERO, targeting individuals between 2 months to 20 years of age residing or attending school in SLSJ. The disease incidence in SLSJ from 2014 to 2018 decreased in the targeted age group (11, 23).

In the UK, a national immunization program (NIP) with BEXSERO was initiated in September 2015 using a two-dose schedule in infants (at 2 and 4 months of age) followed by a booster (at 12 months of age). Public Health England is conducting a 3-year observational study at the national level covering the entire birth cohort (study V72\_38OB).

### **Mechanism of Action**

BEXSERO is a vaccine containing both purified, recombinant protein antigens and OMV derived from *N. meningitidis*. Antigens are variably expressed on meningococci and more than one antigen may be expressed. The inclusion of four antigens, each targeting different steps of meningococcal pathogenesis, may provide multiple targets for vaccine-induced antibodies, and protection against invasive serogroup B strains, even when the expression of one component is low or antigenically different (17, 18). Factor H binding protein (fHbp) is a lipoprotein that binds complement regulatory factor H, thus allowing the evasion of complement-mediated killing and bacterial survival in the blood (12, 13). *Neisseria* adhesin A (NadA) promotes adherence to and invasion of human epithelial cells during meningococcal infection and may also be important for colonization (19, 21, 22). *Neisseria* heparin-binding antigen (NHBA) binds heparin, which may promote bacterial survival in the blood and mediate adhesion to epithelial cells (14, 15). The fourth component are outer membrane vesicles (OMVs) of the epidemic strain from New Zealand, with PorA P1.4 (Porin A serosubtype P1.4) as the immunodominant protein (16).

Protection against IMD is mediated mainly by bactericidal antibodies directed against components of the bacterium. Genes for at least one BEXSERO component have been shown to be present within over 99% of invasive serogroup B strains (20). Immunization with BEXSERO is intended to raise the titer of bactericidal antibodies that specifically bind the vaccine antigens fHbp, NadA, NHBA and PorA P1.4. Meningococci that express either the PorA P1.4 antigen or sufficient levels of any of the other antigens (NadA, fHbp, NHBA), defined as the positive

bactericidal threshold, are predicted to be susceptible to killing by vaccine-elicited immune serum (see Epidemiology Data section).

### **Pharmacodynamics**

No clinical efficacy studies have been undertaken with BEXSERO. Goldschneider *et al.* demonstrated an inverse relationship between meningococcal disease incidence and prevalence of hSBA for serogroups B, C, and A (5, 6). The efficacy of BEXSERO has been inferred by measuring bactericidal antibody responses against hSBA values, the surrogate of protection for serogroup B meningococci (5, 6), to each of the vaccine antigens fHbp, NadA, NHBA and PorA P1.4, using a set of four meningococcal serogroup B reference strains (H44/76, 5/99, M10713 and NZ98/254, respectively). However, data are not available from all vaccine schedules using strain M10713. See Immunogenicity Data section.

Two reference strains (strains H44/76 and 5/99) were selected for hSBA with high level expression of the antigens included in the vaccine, as compared to most of the circulating strains. These two strains could generate a higher percentage of subjects with hSBA  $\geq 1:5$  and higher GMTs than the strains with a low expression (if selected as the reference strains). Although the reference strains are intended to evaluate how well vaccinees mount a functional, antigen-specific immune response against the vaccine antigens, using the strains with high level of antigen expression could potentially result in a more favourable outcome than strains with low level of expression.

### **Pharmacokinetics**

Evaluation of pharmacokinetic properties is not required for vaccines; therefore, no pharmacokinetic studies have been conducted with the vaccine.

## **STORAGE AND STABILITY**

Store in a refrigerator at 2°C to 8°C. Do not freeze. Do not use vaccine that may have been frozen. Protect the vaccine from light.

The expiry date of the vaccine is indicated on the label and packaging. Do not use the vaccine after the expiry date shown on the label.

In the absence of compatibility studies, BEXSERO must not be mixed with other medicinal products.

## **SPECIAL HANDLING INSTRUCTIONS**

A fine off-white deposit may form when the product stands for a long period. Shake the vaccine well before use to form a homogeneous suspension. The vaccine should be visually inspected for particulate matter and discoloration prior to administration.

In the event of any foreign particulate matter and/or variation of physical aspect being observed, do not administer the vaccine.

Any unused product or waste material should be disposed in accordance with local requirements.

## DOSAGE FORMS, COMPOSITION AND PACKAGING

### **Dosage Forms**

BEXSERO is a white opalescent liquid suspension for intramuscular injection.

### **Composition**

1 dose (0.5 mL) contains:

Recombinant <i>Neisseria meningitidis</i> serogroup B NHBA fusion protein <sup>1, 2, 3</sup>	50 micrograms (mcg)
Recombinant <i>Neisseria meningitidis</i> serogroup B NadA protein <sup>1, 2, 3</sup>	50 micrograms (mcg)
Recombinant <i>Neisseria meningitidis</i> serogroup B fHbp fusion protein <sup>1, 2, 3</sup>	50 micrograms (mcg)
Outer membrane vesicles (OMV) from <i>Neisseria meningitidis</i> serogroup B strain NZ98/254 measured as amount of total protein containing the PorA P1.4 <sup>2</sup>	25 micrograms (mcg)

<sup>1</sup> Produced in *E. coli* by recombinant DNA technology.

<sup>2</sup> Adsorbed on aluminium hydroxide (0.5 mg Al<sup>3+</sup>).

<sup>3</sup> NHBA (Neisserial Heparin Binding Antigen), NadA (*Neisseria* adhesin A), fHbp (factor H binding protein).

### **Excipients**

Aluminium hydroxide 1.5 mg, histidine 0.776 mg, sodium chloride 3.125 mg, sucrose 10 mg, water for injections. Residue from the manufacturing process: kanamycin.

### **Packaging**

BEXSERO is supplied as a 0.5 mL suspension in a pre-filled syringe (Type I glass).

The tip cap of the syringe may contain natural rubber latex (see section WARNINGS AND PRECAUTIONS).

Packs of 1 or 10 syringes, supplied with or without needles.

## PART II: SCIENTIFIC INFORMATION

### PHARMACEUTICAL INFORMATION

#### Drug Substance

Proper name: Multicomponent Meningococcal B Vaccine (recombinant, adsorbed)

#### Product Characteristics

BEXSERO is a multicomponent Meningococcal B Vaccine and appears as white opalescent liquid suspension for intramuscular injection in a prefilled syringe. The sequences of the recombinant protein antigens are derived from the following *N. meningitidis* serogroup B strains: NHBA is derived from strain NZ 98/254 and is fused with accessory protein 953 derived from strain 2996; NadA is derived from strain 2996 and fHbp is derived from strain MC58 and is fused with accessory protein 936, derived from strain 2996. The OMV antigen is a suspension that consists of small, membranous spherical vesicles, or fragments of vesicles, in which the native complex antigen composition of the subcapsular cell surface of *N. meningitidis* serogroup B, strain NZ98/254 (B:4:P1.7-2,4) is highly conserved and contains outer membrane protein PorA P1.4 as the main antigen. The recombinant proteins are prepared by recombinant DNA technology using extrachromosomal expression plasmid vectors in *Escherichia coli* cells. The OMV antigen is produced by fermentation of *N. meningitidis* strain NZ98/254, followed by inactivation of the bacteria with deoxycholate, which also mediates vesicle formation.

### CLINICAL TRIALS

#### Study Demographics and Trial Design

Table 14 - Study Demographics and Trial Designs

Study No.	Age at Enrollment	Trial design	Dosage, route of administration and schedule	No. of subjects enrolled (No. receiving BEXSERO <sup>a</sup> )	Mean age of enrolled subjects (range)	Gender of enrolled subjects (% male)
<b>Infants and Children (2 months to 10 years of age)</b>						
V72P12 Phase 2b	2 mos	Open-label, multicenter, randomized, controlled  Safety, immunogenicity, schedule finding	0.5 mL, IM  Schedule: 2, 4, 6 and 2, 3, 4 mos of age	1,885 (1,570)	68.7  (50-107) days	51%

Study No.	Age at Enrollment	Trial design	Dosage, route of administration and schedule	No. of subjects enrolled (No. receiving BEXSERO*)	Mean age of enrolled subjects (range)	Gender of enrolled subjects (% male)
<b>Infants and Children (2 months to 10 years of age)</b>						
V72P12E1 Phase 2b	12, 18, 24 mos	Open-label, multicenter, extension  Safety, immunogenicity of booster in subjects who received a 3-dose series as infants in Study V72P12	0.5 mL, IM Schedule: 4th dose booster at 12, 18, or 24 mos of age; 2 catch up doses at 12, 14 or 18, 20 or 24, 26 mos of age	1,588 (1,519) (1,174: 4th dose; 345: 2-doses in naïve)	17.1 (11-26) mos	52%
V72P13 Phase 3	2 mos	Partially blinded, multicenter, randomized controlled  Safety, immunogenicity, lot consistency	0.5 mL, IM Schedule: 2, 4, 6 mos of age	3,630 (2,480)	73.5 (54-132) days	51%
V72P13E1 Phase 3	12 mos	Open-label, multicenter, randomized  Safety, immunogenicity of fourth dose, 2 catch-up doses starting at 12 or 13 mo for original control group in V72P13	0.5 mL, IM Schedule: 4th dose at 12 mos; 1 or 2 catch-up doses starting at 12 or 13, mos of age	2,249 (2,247) (1,555: 4th dose; 692: 1 or 2 catch-up doses)	12.3 (11-15) mos	51%
V72P13E2 Phase 3	24-27 mos	Open label, randomized, multicenter, extension study	0.5 mL, IM Schedule: 3rd dose boost at 12 mos after 2 catch up doses at 13 and 15 mos or 12 and 14 mos; 2 catch up doses in naïve children (less than 2 years of age) at 24, 26 mos of age	508 (193) (85: 3rd dose; 108: 2-doses in naïve)	25.4 (23-30) mos	52%
V72P16 Phase 2	2 mos	Partially observer-blind, randomized, controlled, multicenter dose-ranging and formulation-finding  Safety and immunogenicity	0.5 mL, IM Schedule: 2, 3, 4, 12 mos of age	1,507 (736)	74.6 (54-91) days	54%

Study No.	Age at Enrollment	Trial design	Dosage, route of administration and schedule	No. of subjects enrolled (No. receiving BEXSERO <sup>®</sup> )	Mean age of enrolled subjects (range)	Gender of enrolled subjects (% male)
<b>Infants and Children (2 months to 10 years of age)</b>						
V72P9 Phase 2	6-8 mos	Single-blind, single center, randomized Safety, immunogenicity of infant primary series + third dose	0.5 mL, IM Schedule: 6, 8, 12 mos of age	60 (30)	7.1 (6-8) mos	47%
V72P9E1 Phase 2	40-60 mos	Open-label, single center, extension Antibody persistence, safety, and immunogenicity of booster dose in children who received a 3-dose series as infants in Study V72P9	0.5 mL, IM Schedule: 4th dose boost at 40 mos; 2 catch up doses in naive children at 40,42 or 60,62 mos	120 (103) (14: 4th dose; 89: 2-doses in naive)	50.0 (39-62) mos	48%
V72P6E1 Phase 2	40-60 mos	Open-Label, single center, extension Antibody persistence, safety and immunogenicity of booster doses in children who received 1 or 4 doses as infants in study V72P6	0.5 mL, IM Schedule: 5th dose boost at 40 mos; 2 doses at 40, 42 mos after one dose at 12 mos; 2 catch up doses in naive children at 40, 42 mos.	113 (69) (19: 5th dose; 8: 2-doses after one dose at 12 mos; 42: 2-doses in naive)	41.5 (40-44) mos	50%
V72_28 Phase 3	2 mos-8 yrs	Open-label, multicenter, safety, tolerability and immunogenicity	0.5 mL, IM Schedules: Group I <sup>b</sup> : 2½, 3½, 5 + booster at 11 months of age Group II <sup>b</sup> : 3½, 5 + booster at 11 months of age Group III <sup>b</sup> : 6, 8 mos + booster at 11 months of age Group IV <sup>c</sup> : 0, 2 mos Group V <sup>d</sup> : 3, 5, 7, 12 mos Group VI <sup>d</sup> : 3, 5, 7, 12, 13, 15 mos	Group I: 253 (252) Group II: 250 (249) Group III: 251 (250) Group IV: 404 (404) Group V: 126 (126) Group VI: 125 (124)	Group I: 2 mos Group II: 3 mos Group III: 6 mos Group IV: 7 yrs Group V: 3 mos Group VI: 3 mos	Group I: 54% Group II: 50% Group III: 49% Group IV: 51% Group V: 41% Group VI: 53%

Study No.	Age at Enrollment	Trial design	Dosage, route of administration and schedule	No. of subjects enrolled (No. receiving BEXSERO®)	Mean age of enrolled subjects (range)	Gender of enrolled subjects (% male)
<b>Infants and Children (2 months to 10 years of age)</b>						
V72_28E1 Phase 3	35 mos – 12 yrs	Open label, multicenter extension study of Study V72_28	0.5 mL, IM Group I-III: Vaccinated cohort: subjects from Groups I-III of Study V72_28 receiving 1 dose 24-36 mos later.  Group IVa and IVb: Vaccinated cohort: subjects from Group IV of Study V72_28 receiving 1 dose 24-36 mos later.	Group I vaccinated: 98 (97) Group 1 non-vaccinated: 47 (n.a)  Group II vaccinated: 89 (89), non-vaccinated: 43 (n.a)  Group III, vaccinated: 81 (80); non-vaccinated: 39 (n.a)  Group IVa: vaccinated: 32 (32); non-vaccinated 36 (n.a)  Group IVb, vaccinated: 91 (91); non-vaccinated: 90 (n.a)	Groups I-III: 35-47 months of age  Group IVa: approximate age range 4-7 years of age (i.e 24-36 months after Study V72_28)  Group IVb: approximate age range 8-12 years of age (i.e 24-36 months after Study V72_28)	Group I vaccinated: 55% Group 1 non-vaccinated: 43%  Group II vaccinated: 46%, non-vaccinated: 49%  Group III, vaccinated: 57%; non-vaccinated: 41%  Group IVa, vaccinated: 41%; non-vaccinated 58%  Group IVb, vaccinated: 52%; non-vaccinated: 49%
V72_56 Phase 3	3 months	Open-label, randomized, multicenter, non-inferiority	0.5 mL, IM Schedule: 3, 5, 7 and 13 mos	750 (744)	102 (91-113) days	n/a
V72_62 Phase 3	2-17 yrs	Open-label, controlled, multicenter, safety, tolerability and immunogenicity	0.5 mL, IM Schedule: 0, 2 mos	239 (n.a.)	10.3 (2-17) yrs	55%
<b>Adolescents and Adults (11 years through 25 years of age)</b>						
V72P10 Phase 2b/3	11-17 yrs	Observer-blind, multicenter, randomized, placebo controlled  Safety, immunogenicity, schedule finding	0.5 mL, IM Schedule: mos 0; mos 6; mos 0, 1; mos 0, 2; mos 0, 6; mos 0, 1, 2; mos 0, 1, 6; mos 0, 2, 6	1,631 (1,622)	13.8 (10-17) yrs	44%



Study No.	Age at Enrollment	Trial design	Dosage, route of administration and schedule	No. of subjects enrolled (No. receiving BEXSERO <sup>a</sup> )	Mean age of enrolled subjects (range)	Gender of enrolled subjects (% male)
<b>Adolescents and Adults (11 years through 25 years of age)</b>						
V72_41 Phase 3	11-17 yrs	Observer-blind, multicenter, randomized Safety and immunogenicity, lot consistency	0.5 mL, IM Schedule: 0, 1 mos	344 (342)	13.7 (11-17) yrs	55%
V72_75 Phase 3b	15-24 yrs	Open label, multicenter, controlled, extension study of V72P10 and V72_41  Immunogenicity, antibody persistence in adolescents and young adults	0.5 mL, IM Schedule: 0, 1; 0,2; 0,6 mos	531 (530) Follow-on subjects:275 Vaccine Naïve subjects: 255	19.7 (15-24) yrs	51%

<sup>a</sup> Number of subjects in the BEXSERO safety population. Defined as those subjects who were vaccinated with BEXSERO and who provided some post-baseline safety data;

<sup>b</sup> Groups I, II and III: Subjects Receiving a 2-or 3-Dose Primary Series Followed by a Booster Dose

<sup>c</sup> Group IV: Subjects Receiving a 2-Dose Catch-Up Series, Group IVa (children 2-5 years of age); Group IVb (children 6-10 years of age)

<sup>d</sup> Groups V and VI: Infants 3 months of age receiving a 2-dose primary series followed by a booster dose of BEXSERO with concomitant MenC-CRM or MenC-CRM alone

IM: intramuscular;

mos: months; yrs: years.

### Immunogenicity Data

Immunogenicity of BEXSERO was evaluated in primary and extension studies. Most of the primary immunogenicity studies were conducted as randomized, multicenter clinical trials that enrolled infants, children, adolescents and adults (see Table 14). The primary immunogenicity measure was the proportion of subjects with human serum bactericidal assay (hSBA) equal to or above the threshold of 1:4 against each of the meningococcal serogroup B reference strains. This threshold, used in early-stage clinical studies (V72P6, V72P9, V72P4, V72P5 and V72P10) and in their extensions (V72P6E1, V72P9E1, V72P10E1), is an accepted correlate of protection. A threshold of 1:5 was then set after hSBA assay validation to ensure, based on the intermediate precision of the assay, 95% certainty of a true response of 1:4, and this cutoff was used to define seropositive responses in late-stage clinical studies.

**Table 15 - Summary of the Main Studies**

Study	Primary Immunogenicity Objectives	Prospectively Defined Criterion	Outcome
<p><b>V72P12</b> <b>Phase 2b</b></p>	<ul style="list-style-type: none"> <li>Demonstration of a sufficient immune response to BEXSERO when given concomitantly with routine vaccines to healthy infants at either 2, 4 and 6 or 2, 3 and 4 months of age, by evaluation of hSBA at 1 month after the third vaccination.</li> </ul>	<ul style="list-style-type: none"> <li>The immune response was considered sufficient if the lower limit of the two-sided 95% CI for the percentage of subjects with hSBA <math>\geq 1:5</math> at 1 month following the third vaccination was <math>\geq 70\%</math> for all 3 reference strains H44/76, NZ98/254 and 5/99.</li> </ul>	<p>Objective was met.</p> <ul style="list-style-type: none"> <li>The lower limits of the two-sided 95% CI for the percentage of subjects with hSBA <math>\geq 1:5</math> for the 2, 4, 6-month schedule were: 98% for strain H44/76, 98% for strain 5/99 and 75% for strain NZ98/254.</li> <li>The lower limits of the two-sided 95% CI for the percentage of subjects with hSBA <math>\geq 1:5</math> for the 2,3,4-month schedule were: 97% for strain H44/76, 99% for strain 5/99 and 76% for strain NZ98/254.</li> </ul>
<p><b>V72P13</b> <b>Phase 3</b></p>	<p>Two Co-primary Objectives:</p> <ul style="list-style-type: none"> <li>To show the consistency of the immune response from 3 lots of BEXSERO, by hSBA GMTs, when administered to healthy infants at 2, 4 and 6 months of age, at 1 month after the third vaccination.</li> <li>Demonstration of a sufficient immune response to BEXSERO (3 lots combined) when given concomitantly with routine vaccines to healthy infants at 2, 4 and 6 months of age, by evaluation of hSBA at 1 month after the third vaccination.</li> </ul>	<ul style="list-style-type: none"> <li>The 3 BEXSERO vaccine lots were considered equivalent if for each of the reference strains H44/76, NZ98/254 and 5/99 and each pair of vaccine lots, the two-sided 95% CI of the ratio of GMTs at 1 month after the third vaccination was contained within the interval [0.50, 2.00].</li> <li>The immune response was considered sufficient immune if the lower limit of the two-sided 95% CI for the percentage of subjects with hSBA <math>\geq 1:5</math> at 1 month following the third vaccination was <math>\geq 70\%</math> for all 3 reference strains H44/76, NZ98/254 and 5/99, for the 3 BEXSERO lots combined.</li> </ul>	<p>Objectives were met.</p> <ul style="list-style-type: none"> <li>For each reference strain, for all 3 pairs of BEXSERO vaccine lots simultaneously, the two-sided 95% CI for the ratios of GMTs at 1 month after the third vaccination were entirely contained within the interval [0.74, 1.33], thereby meeting the criterion for lot consistency [0.50, 2.00].</li> <li>The lower limits of the two-sided 95% CI for the percentage of subjects with an hSBA <math>\geq 1:5</math> at 1 month following the third vaccination were: 100% against the H44/76 and 5/99 strains, and 84% against the NZ98/254 strain, thereby meeting the sufficient immune response criterion.</li> </ul>
<p><b>V72P13E1</b> <b>Phase 3</b></p>	<ul style="list-style-type: none"> <li>Demonstration of a sufficient immune response following a fourth dose of BEXSERO administered at 12 months of age, either with or without concomitant Priorix-Tetra vaccination, to children (less than 2 years of age) previously primed with three doses of BEXSERO as infants in Study V72P13.</li> </ul>	<ul style="list-style-type: none"> <li>The fourth dose immune response was considered sufficient if for the percentage of subjects with hSBA <math>\geq 1:5</math>, the lower limit of the two-sided 95% CI was <math>\geq 75\%</math> for all 3 reference strains H44/76, NZ98/254 and 5/99.</li> </ul>	<p>Objectives were met.</p> <ul style="list-style-type: none"> <li>For strains H44/76 and 5/99, 100% of the subjects had hSBA <math>\geq 1:5</math>. The lower limit of the two-sided 95% CI was 98% in subjects with or without concomitant Priorix-Tetra vaccination.</li> <li>For strain NZ98/254, 97% and 94% of the subjects in the vaccination groups had hSBA <math>\geq 1:5</math>. The lower limit of the two-sided 95% CI was 93% in subjects with concomitant Priorix-Tetra and 90% in subjects without concomitant Priorix-Tetra vaccination.</li> </ul>

Study	Primary Immunogenicity Objectives	Prospectively Defined Criterion	Outcome
V72_41 Phase 3	<ul style="list-style-type: none"> <li>Demonstration of the equivalence of rMenB+OMV NZ lot 1 to rMenB+OMV NZ lot 2 when administered to adolescents, as measured by hSBA GMTs for strains H44/76, 5/99, and NZ98/254 and ELISA GMCs against vaccine antigen 287-953 approximately 30 days after a primary vaccination course of two doses administered one month apart.</li> </ul>	<ul style="list-style-type: none"> <li>The equivalence was considered a success if, at one month following the second vaccination, the two-sided 95% confidence interval (CI) of the ratio of the hSBA GMTs for each of 3 serogroup B reference strains (H44/76, 5/99, and NZ98/254) and the two-sided 95% CI of the ratio of the ELISA GMCs against vaccine antigen 287-953 are contained within the interval (0.5, 2.0).</li> </ul>	<p>Objective was met.</p> <ul style="list-style-type: none"> <li>The ratios of hSBA GMTs in Lot 1_Rosia to Lot 2_Siena at one month after the second vaccination were 1.0, 0.92, and 0.81 for strains H44/76, 5/99, and NZ98/254, respectively, with corresponding two-sided 95% confidence intervals of (0.82, 1.23), (0.77, 1.10), and (0.60, 1.09).</li> <li>The ratio of ELISA GMCs against vaccine antigen 287-953 at one month after second vaccination was 0.83, with a corresponding two-sided 95% CI of (0.67, 1.02).</li> </ul>

CI: confidence interval;

GMT: geometric mean titers

hSBA: serum bactericidal assay using human complement

### **Immunogenicity in Infants aged 2 to 5 months**

#### ***3+1 Infant Schedule***

Immunogenicity results at one month after three doses of BEXSERO administered at 2, 3, 4 and 2, 4, 6 months of age are summarized in Table 16. Persistence and data after a fourth dose administered at 12 months of age (following administration at 2, 3, 4 months of age in Study V72P12E1 and at 2, 4, 6 months of age in Study V72P13E1) are summarized in Table 17.

Baseline Geometric Mean Titers (GMT) were uniformly low against all strains in the BEXSERO (ranging from 1.02 to 1.49 for fHbp, NadA and PorA P1.4 antigens and from 3.15 to 3.51 for NHBA) and the control groups (ranging from 1.01 to 1.28 for fHbp, NadA and PorA P1.4 antigens and was 3.91 for NHBA) across studies. The responses one month after the third vaccination at a 2, 4, 6-month schedule were high against all antigens in the BEXSERO groups (Table 16). In contrast, the mean hSBA GMTs remained low and similar with respect to the baseline in the control groups (ranging from 1.04 to 1.25).

**Table 16 - Serum Bactericidal Antibody Responses at 1 Month Following the Third Dose of BEXSERO given at 2, 3, 4 or 2, 4, 6 Months of Age (V72P12, V72P13 and V72P16)**

Antigen		Study V72P13 2, 4, 6 months	Study V72P12 2, 3, 4 months	Study V72P16 2, 3, 4 months
fHbp	% seropositive <sup>a</sup> (95% CI)	N=1,149 100% (99-100)	N=273 99% (97-100)	N=170 100% (98-100)
	hSBA GMT (95% CI)	91 (87-95)	82 (75-91)	101 (90-113)
NadA	% seropositive <sup>a</sup> (95% CI)	N=1,152 100% (99-100)	N=275 100% (99-100)	N=165 99% (97-100)
	hSBA GMT (95% CI)	635 (606-665)	325 (292-362)	396 (348-450)
PorA P1.4	% seropositive <sup>a</sup> (95% CI)	N=1,152 84% (82-86)	N=274 81% (76-86)	N=171 78% (71-84)
	hSBA GMT (95% CI)	14 (13-15)	11 (9.14-12)	10 (8.59-12)
NHBA	% seropositive <sup>a</sup> (95% CI)	N=100 84% (75-91)	N=112 37% (28-46)	N=35 43% (26-61)
	hSBA GMT (95% CI)	16 (13-21)	3.24 (2.49-4.21)	3.29 (1.85-5.83)

<sup>a</sup> % seropositive = the percentage of subjects who achieved an hSBA  $\geq$  1:5

hSBA = Serum Bactericidal Assay using human complement

GMT = Geometric Mean Titer.

N= number of subjects with evaluable serum for analysis of that strain.

As compared with study V72P13 (2, 4, 6- month schedule), percentages of subjects with hSBA  $\geq$ 1:5 against NHBA and GMTs against NHBA, NadA and PorA P1.4 were significantly lower in study V72P12 and V72P16 (2, 3, 4- month schedule) at one month after the third vaccination.

A modest response was demonstrated following vaccinations with BEXSERO at the 2, 3, 4- month schedule in studies V72P12 and V72P16 as the percentages of subjects with hSBA  $\geq$ 1:5 against NHBA was 36% vs. 6%; 43% vs. 20% for the BEXSERO vs. control groups, respectively.

### **2+1 Infant Schedule**

The immunogenicity after two doses (at 3½ and 5 months of age) or three doses (at 2½, 3½ and 5 months of age) of BEXSERO followed by a booster has been evaluated in an additional phase 3 clinical study (Study V72\_28). The percentages of seropositive subjects (i.e. achieving an hSBA of at least 1:4) ranged from 44% to 100% one month after the second dose and from 55% to 100% one month after the third dose, respectively for the two-dose schedule and the three-dose schedule. At one month following a booster administered 6 months after the last primary dose, the percentages of seropositive subjects ranged from 87% to 100% for the two-dose schedule, and from 83% to 100% for the three-dose schedule.

***Antibody Persistence following 2+1 or 3+1 Infant Schedule***

The antibodies against PorA and fHbp rapidly declined in infants 6 and 12 months after the third dose, respectively. However, a booster response was observed following a fourth vaccine dose administered during the second year of life, consistent with adequate priming with a three-dose primary series (Table 17).

Antibody persistence was evaluated in an extension study in children 3 to 4 years of age (Study V72\_28E1). At 2 to 3 years after the completion of the vaccination course in Study V72\_28, the antibody levels declined against all strains in both 2+1 and 3+1 schedules; the percentages of seropositive subjects (i.e. achieving an hSBA of at least 1:4) ranged from 35% to 91% or from 36% to 84%, respectively. In the same study the response to an additional dose administered 2 to 3 years after the booster was indicative of immunological memory as shown by a robust antibody response against all BEXSERO antigens, with seropositive rates ranging from 81% to 100% and from 70% to 99%, respectively.

**Table 17 - Serum Bactericidal Antibody Responses Following a Booster at 12 Months After a Primary Series Administered at 2, 3 and 4 or 2, 4 and 6 Months of Age, and Persistence of Bactericidal Antibody One Year After the Booster (V72P12E1 and V72P13E1)**

Antigen		2, 3, 4, 12 months	2, 4, 6, 12 months
<b>fHbp</b>	pre-booster <sup>a</sup>	N=81	N=426
	% seropositive <sup>b</sup> (95% CI)	58% (47-69)	82% (78-85)
	hSBA GMT (95% CI)	5.79 (4.54-7.39)	10 (9.55-12)
	1 month after booster	N=83	N=422
	% seropositive <sup>b</sup> (95% CI)	100% (96-100)	100% (99-100)
	hSBA GMT (95% CI)	135 (108-170)	128 (118-139)
<b>NadA</b>	pre-booster <sup>a</sup>	N=79	N=423
	% seropositive <sup>b</sup> (95% CI)	97% (91-100)	99% (97-100)
	hSBA GMT (95% CI)	63 (49-83)	81 (74-89)
	1 month after booster	N=84	N=421
	% seropositive <sup>b</sup> (95% CI)	100% (96-100)	100% (99-100)
	hSBA GMT (95% CI)	1558 (1262-1923)	1465 (1350-1590)
<b>PorA P1.4</b>	pre-booster <sup>a</sup>	N=83	N=426
	% seropositive <sup>b</sup> (95% CI)	19% (11-29)	22% (18-26)
	hSBA GMT (95% CI)	1.61 (1.32-1.96)	2.14 (1.94-2.36)
	1 month after booster	N=86	N=424
	% seropositive <sup>b</sup> (95% CI)	97% (90-99)	95% (93-97)
	hSBA GMT (95% CI)	47 (36-62)	35 (31-39)
<b>NHBA</b>	pre-booster <sup>a</sup>	N=69	N=100
	% seropositive <sup>b</sup> (95% CI)	25% (15-36)	61% (51-71)
	hSBA GMT (95% CI)	2.36 (1.75-3.18)	8.4 (6.4-11)
	1 month after booster	N=67	N=100
	% seropositive <sup>b</sup> (95% CI)	76% (64-86)	98% (93-100)
	hSBA GMT (95% CI)	12 (8.52-17)	42 (36-50)
<b>NHBA</b>	12 months after booster	N=291	N=291
	% seropositive <sup>b</sup> (95% CI)	-	36% (31-42%)
	hSBA GMT (95% CI)	-	3.35 (2.88-3.9)

<sup>a</sup> pre-booster time point represents persistence of bactericidal antibody at 8 months after BEXSERO vaccination at 2, 3 and 4 months of age and 6 months after BEXSERO vaccination at 2, 4 and 6 months of age.

<sup>b</sup> % seropositive = the percentage of subjects who achieved an hSBA  $\geq$  1:5

hSBA = Serum Bactericidal Assay using human complement; GMT = Geometric Mean Titer

N= number of subjects with evaluable serum for analysis of that strain.

### **Concomitant use of BEXSERO with routine vaccines**

Concomitant administration of BEXSERO was studied with any of the following vaccine antigens, either as monovalent or as combination vaccines: diphtheria, tetanus, acellular pertussis, *Haemophilus influenzae* type b, inactivated poliomyelitis, hepatitis B, heptavalent pneumococcal conjugate, measles, mumps, rubella, varicella, and meningococcal groups A, C, W, Y conjugate.

Clinical study V72P12 demonstrated that the percentage of subjects with hSBA  $\geq 1:5$  for strain NZ98/254 was lower in the group that concomitantly administered BEXSERO and the routine vaccines (combined DTaP-IPV-HBV/Hib vaccine and heptavalent pneumococcal conjugate vaccine) than the group where they were administered separately at 1 month after the third dose. When administered alone, BEXSERO also elicited higher hSBA GMTs for all strains as compared to the concomitant group. The clinical implication of these differences remains unknown.

Inconsistent results were seen across studies for responses to inactivated poliovirus type 2 and pneumococcal conjugate serotype 6B; lower antibody titers to the pertussis pertactin antigen were also noted.

A clinical study for concomitant use of BEXSERO with monovalent MenC-CRM conjugate vaccine showed that one month following 2 doses vaccination course and one month post-booster dose, the percentages of subjects achieving hSBA titers  $\geq 8$  against meningococcal serogroup C were comparable in both groups. Lower geometric mean titers (GMTs) were observed for meningococcal serotype-C (MenC-CRM vaccine) in the co-administration group than those seen in MenC-CRM administered alone. Given that the titers in both groups were high, it is not likely that this difference will have a clinically significant impact.

A clinical study for concomitant use of BEXSERO with combined Men A, C, W, Y conjugate vaccine showed a robust immune response one month following 3 doses vaccination course and a booster, indicating that the concomitant administration is non-inferior to that of either vaccine administered alone.

In addition, concomitant use of BEXSERO and MMRV demonstrated non-inferiority of seroconversion ( $\geq 1.25$  gpELISA units/mL), but not of seroprotection ( $\geq 5$  gpELISA units/mL) for varicella after the first dose, although the difference between the groups was only 2% (95% CI, -11%, 7%). The clinical implication of these differences remains unknown.

In a clinical trial, prophylactic use of acetaminophen had no impact on the immune responses of BEXSERO and for most antigens in routine vaccines after the primary series. These data do not suggest any clinically significant interference also considering that no impact was observed on the immune responses after the booster doses.

### **Immunogenicity in Infants aged 6 to 11 months, Children aged 12 months to 10 years**

The immunogenicity after two doses in infants and children has been documented in three studies whose results are summarized in Table 18. Baseline GMTs were uniformly low against all three strains in the studies in infants 6 to 11 months of age and children 12 to 23 months of age (ranging from 1.00 to 1.70) and increased following vaccination. The increase in hSBA titers for vaccine antigens was similar in additional groups of children following BEXSERO

vaccination at 12-14 and 18-20 months of age. In these additional groups a similar response was also observed in terms of percentages of seropositive subjects (100% against fHbp antigen; 98-100% against NadA antigen; 93-99% against PorA P1.4 antigen; 74-86% against NHBA antigen).

In 24 to 26 months old children, baseline GMTs were also low (ranging from 1.01 to 2.32 across all vaccine antigens). Additional data relative to the administration of two BEXSERO doses 2 months apart in children at 40-42 and 60-62 months of age (Studies V72P6E1, V72P9E1) were in line with the responses presented in Table 18.

**Table 18 - Serum Bactericidal Antibody Responses Following BEXSERO Vaccination at 6 and 8 Months of Age (V72P9), 13 and 15 Months of Age (V72P13E1) or 24 and 26 Months of Age (V72P13E2) and Persistence of Bactericidal Antibody One Year After the Two Doses at 13 and 15 Months of Age (V72P13E2)**

Antigen		Age range		
		6 to 11 months of age	12 to 23 months of age	24 to 30 months of age
		Age of vaccination		
		6, 8 months	13, 15 months	24, 26 months
fHbp	<u>1 month after 2<sup>nd</sup> dose</u> % seropositive <sup>a</sup> hSBA GMT	N=23 100% 250	N=163 100% (98-100) <sup>b</sup> 271 (237-310) <sup>b</sup>	N=105 100% (97-100) <sup>b</sup> 220 (186-261) <sup>b</sup>
	<u>12 months after 2<sup>nd</sup> dose</u> % seropositive <sup>a</sup> hSBA GMT	-	N=68 74% (61-83) <sup>b</sup> 14 (9.4-20) <sup>b</sup>	-
NadA	<u>1 month after 2<sup>nd</sup> dose</u> % seropositive <sup>a</sup> hSBA GMT	N=23 100% 534	N=164 100% (98-100) <sup>b</sup> 599 (520-690) <sup>b</sup>	N=103 99% (95-100) <sup>b</sup> 455 (372-556) <sup>b</sup>
	<u>12 months after 2<sup>nd</sup> dose</u> % seropositive <sup>a</sup> hSBA GMT	-	N=68 97% (90-100) <sup>b</sup> 70 (47-104) <sup>b</sup>	-
PorA P1.4	<u>1 month after 2<sup>nd</sup> dose</u> % seropositive <sup>a</sup> hSBA GMT	N=22 95% 27	N=164 100% (98-100) <sup>b</sup> 43 (38-49) <sup>b</sup>	N=108 98% (93-100) <sup>b</sup> 27 (23-32) <sup>b</sup>
	<u>12 months after 2<sup>nd</sup> dose</u> % seropositive <sup>a</sup> hSBA GMT	-	N=68 18% (9-29) <sup>b</sup> 1.65 (1.2-2.28) <sup>b</sup>	-
NHBA	<u>1 month after 2<sup>nd</sup> dose</u> % seropositive <sup>a</sup> hSBA GMT	-	N=46 63% (48-77) <sup>b</sup> 11 (7.07-16) <sup>b</sup>	N=100 97% (91-99) <sup>b</sup> 38 (32-45) <sup>b</sup>
	<u>12 months after 2<sup>nd</sup> dose</u> % seropositive <sup>a</sup> hSBA GMT	-	N=65 38% (27-51) <sup>b</sup> 3.7 (2.15-6.35) <sup>b</sup>	-

<sup>a</sup> % seropositive = the percentage of subjects who achieved an hSBA  $\geq$  1:4 (in the 6 to 11 months range of age) and hSBA  $\geq$  1:5 (in the 12 to 23 months range).

<sup>b</sup> 95% Confidence Intervals are reported in brackets only for data generated from clinical studies V72P13E1 and V72P13E2 in the age range 12 to 23 months.

hSBA = Serum Bactericidal Assay using human complement

GMT = Geometric Mean Titer.



N= number of subjects with evaluable serum for analysis of that strain.

In Study V72\_28, participants received two doses of BEXSERO administered two months apart. The seroresponse rates and hSBA GMTs are shown in Table 19.

**Table 19 - Serum bactericidal antibody responses at 1 month following the second dose of BEXSERO given to children 2-10 years of age following a 0, 2-month schedule (V72\_28)**

Antigen		2 to 5 years of age	6 to 10 years of age
fHbp	% seropositive* (95% CI)	N=99 100% (96-100)	N=287 99% (96-100)
	hSBA GMT** (95% CI)	140 (112-175)	112 (96-130)
NadA	% seropositive (95% CI)	N=99 99% (95-100)	N=291 100% (98-100)
	hSBA GMT (95% CI)	584 (466-733)	457 (392-531)
PorA P1.4	% seropositive (95% CI)	N=100 98% (93-100)	N=289 99% (98-100)
	hSBA GMT (95% CI)	42 (33-55)	40 (34-48)
NHBA	% seropositive (95% CI)	N=95 91% (83-96)	N=275 95% (92-97)
	hSBA GMT (95% CI)	23 (18-30)	35 (29-41)

\* % seropositive = the percentage of subjects who achieved an hSBA  $\geq$  1:4 (against reference strains for fHbp, NadA, PorA P1.4 antigens) and an hSBA  $\geq$  1:5 (against reference strain for NHBA antigen).

\*\* GMT = geometric mean titre.

N= number of subjects with evaluable serum for analysis of that strain.

In the second study (V72\_28E1), two doses of BEXSERO were administered one month apart in children 3 to 12 years of age. The percentages of seropositive children up to 10 years of age (i.e. achieving an hSBA of at least 1:4) across strains ranged from 46% to 95% at one month after the first dose and from 69% to 100% at one month after the second dose.

This study also evaluated antibody persistence and the response to a booster dose in children who received the two-dose primary series at 2-5 or 6-10 years of age. After 24 to 36 months, the percentages of seropositive subjects (i.e., achieving an hSBA of at least 1:4) declined, ranging across strains from 21% to 74% in children 4 to 7 years of age and from 47% to 86% in children 8 to 12 years of age. The response to a booster dose administered 24 to 36 months after the primary series was indicative of immunological memory as the percentages of seropositive subjects ranged across strains from 93% to 100% in children 4 to 7 years of age and from 96% to 100% in children 8 to 12 years of age.

**Immunogenicity in Adolescents aged 11 to 17 years**

The immunogenicity data of two doses administered with an interval of one, two or six months in adolescents (V72P10, V72\_41) are shown in Table 20. Baseline GMTs ranged from 2.61 to 4.11 in adolescents against fHbp, NadA and PorA P1.4 antigens. Baseline GMTs against NHBA antigen ranged from 30 to 32 in adolescents (V72P10).

**Table 20 - Serum Bactericidal Antibody Responses in Adolescents One Month After Two Doses of BEXSERO Administered According to Different Two-Dose Schedules (V72\_41 and V72P10)**

Antigen		V72_41 0, 1 months	V72P10 0, 1 months	V72P10 0, 2 months	V72P10 0, 6 months
fHbp	% seropositive <sup>a</sup>	N=298 99% (98-100) <sup>b</sup>	N=638 100% (99-100) <sup>b</sup>	N=319 100% (99-100) <sup>b</sup>	N=86 100% (99-100) <sup>b</sup>
	hSBA GMT	117 (105-130) <sup>b</sup>	210 (193-229) <sup>b</sup>	234 (209-263) <sup>b</sup>	218 (157-302) <sup>b</sup>
NadA	% seropositive <sup>a</sup>	N=299 100% (99-100) <sup>b</sup>	N=639 100% (99-100) <sup>b</sup>	N=320 99% (98-100) <sup>b</sup>	N=86 99% (94-100) <sup>b</sup>
	hSBA GMT	179 (163-197) <sup>b</sup>	490 (455-528) <sup>b</sup>	734 (653-825) <sup>b</sup>	880 (675-1147) <sup>b</sup>
PorA P1.4	% seropositive <sup>a</sup>	N=298 75% (70-80) <sup>b</sup>	N=639 100% (99-100) <sup>b</sup>	N=319 100% (99-100) <sup>b</sup>	N=86 100% (96-100) <sup>b</sup>
	hSBA GMT	10 (8.77-12) <sup>b</sup>	92 (84-102) <sup>b</sup>	123 (107-142) <sup>b</sup>	140 (101-195) <sup>b</sup>
NHBA	% seropositive <sup>a</sup>	-	N=46 100% (92-100) <sup>b</sup>	N=46 100% (92-100) <sup>b</sup>	-
	hSBA GMT	-	99 (76-129) <sup>b</sup>	107 (82-140) <sup>b</sup>	-

<sup>a</sup> % seropositive = the percentage of subjects who achieved an hSBA  $\geq$  1:4 (in clinical study V72P10) and hSBA  $\geq$  1:5 (in clinical study V72\_41).

<sup>b</sup> 95% Confidence Intervals are reported in brackets only for clinical studies V72P10 and V72\_41.

hSBA = Serum Bactericidal Assay using human complement.

GMT = Geometric Mean Titer.

N= number of subjects with evaluable serum for analysis of that strain.

In study V72P10, subjects were stratified by pre-vaccination titer baseline hSBA <1:4 or ≥1:4. The percentage of subjects with at least a 4-fold increase in hSBA titer from baseline to one month after the last dose of BEXSERO is summarized in Table 21.

**Table 21 - Percentage of Adolescents With Seroreponse and at Least 4-Fold Rise<sup>b</sup> in Bactericidal Titers One Month After Two Doses of BEXSERO Administered According to Different Two-Dose Schedules - Stratified by Pre-Vaccination Titers (V72P10)**

Antigen			0, 1 months	0, 2 months	0, 6 months
<b>fHbp</b>	% seropositive <sup>a</sup> (95% CI)	pre-vaccination titer <1:4	N=369 100% (98-100)	N=179 100% (98-100)	N=55 100% (94-100)
		pre-vaccination titer ≥1:4	N=269 100% (99-100)	N=140 100% (97-100)	N=31 100% (89-100)
	% 4-fold increase (95% CI)	pre-vaccination titer <1:4	N=369 100% (98-100)	N=179 100% (98-100)	N=55 100% (94-100)
		pre-vaccination titer ≥1:4	N=268 90% (86-93)	N=140 86% (80-92)	N=31 90% (74-98)
<b>NadA</b>	% seropositive <sup>a</sup> (95% CI)	pre-vaccination titer <1:4	N=427 100% (99-100)	N=211 99% (97-100)	N=64 98% (92-100)
		pre-vaccination titer ≥1:4	N=212 100% (98-100)	N=109 100% (97-100)	N=22 100% (85-100)
	% 4-fold increase (95% CI)	pre-vaccination titer <1:4	N=426 99% (98-100)	N=211 99% (97-100)	N=64 98% (92-100)
		pre-vaccination titer ≥1:4	N=212 96% (93-98)	N=109 95% (90-98)	N=22 95% (77-100)
<b>PorA P1.4</b>	% seropositive <sup>a</sup> (95% CI)	pre-vaccination titer <1:4	N=427 100% (98-100)	N=208 100% (98-100)	N=64 100% (94-100)
		pre-vaccination titer ≥1:4	N=212 100% (98-100)	N=111 100% (97-100)	N=22 100% (85-100)
	% 4-fold increase (95% CI)	pre-vaccination titer <1:4	N=426 99% (98-100)	N=208 100% (98-100)	N=64 100% (94-100)
		pre-vaccination titer ≥1:4	N=211 81% (75-86)	N=111 77% (68-84)	N=22 82% (60-95)
<b>NHBA</b>	% seropositive <sup>a</sup> (95% CI)	pre-vaccination titer <1:4	N=2 100% (16-100)	N=9 100% (66-100)	-
		pre-vaccination titer ≥1:4	N=44 100% (92-100)	N=37 100% (91-100)	-
	% 4-fold increase (95% CI)	pre-vaccination titer <1:4	N=2 100% (16-100)	N=9 89% (52-100)	-
		pre-vaccination titer ≥1:4	N=44 30% (17-45)	N=37 19% (8-35)	-

<sup>a</sup> % seropositive = the percentage of subjects who achieved an hSBA ≥ 1:4

<sup>b</sup> At least 4-fold rise = a post-vaccination hSBA ≥1:8 for subjects with pre-vaccination hSBA <1:2; a post-vaccination 4-fold rise for subjects with pre-vaccination hSBA ≥1:2.

N= number of subjects with evaluable serum for analysis of that strain.

***Immunogenicity in children and adolescents with complement deficiencies, and in individuals with splenic dysfunction or asplenia***

In a phase 3b, open label clinical study (study V72\_62), children and adolescents 2 to 17 years of age with complement deficiencies (n=40), with asplenia or splenic dysfunction (n=107), and age-matched healthy subjects (n=85) received two doses of BEXSERO two months apart. At 1 month following the 2-dose vaccination course, the percentages of subjects with hSBA  $\geq$  1:5 in individuals with complement deficiencies and asplenia or splenic dysfunction were 87% (95% CI: 72.6-95.7) and 97% (95% CI: 91.8-99.4) for antigen fHbp, 95% (95% CI: 82.3-99.4) and 100% (95% CI: 96.6-100) for antigen NadA, 68% (95% CI: 51.3-82.5) and 86% (95% CI: 77.7-91.9) for antigen PorA P1.4, 73% (95% CI: 55.9-86.2) and 94% (95% CI: 87.8-97.8) for antigen NHBA, respectively, indicating an immune response in these immunocompromised subjects. The percentages of healthy subjects with hSBA  $\geq$  1:5 were 98% (95% CI: 91.8-99.71) for antigen fHbp, 99% (95% CI: 93.5-99.97) for antigen NadA, 83% (95% CI: 73.6-90.6) for antigen PorA P1.4, and 99% (95% CI: 93.5-99.97) for antigen NHBA.

**Adults**

Data from study V72\_75 were obtained in Canada and Australia after two doses of BEXSERO administered according to a 0-1 month schedule. Immunogenicity responses as given by percentages of subjects with hSBA  $\geq$  1:5 (seropositive threshold) and GMTs 1 month following 2 doses (0,1 month) of BEXSERO in subjects 18 to 24 years of age are presented in Table 22. The percentages of subjects with at least a 4-fold rise at one month after the second dose compared to baseline ranged from 24% to 100% for fHbp, NadA and NHBA.

**Table 22 - Serum Bactericidal Antibody Response<sup>a</sup> at 1 month after a 2-Dose Schedule in Canadian and Australian Adults (18 to 24 years of age) (V72\_75)**

Antigen	Study	V72_75 <sup>b</sup>	
		Baseline	1 month after a 2-Dose Schedule (0,1 month)
fHbp	% seropositive (95% CI)	6 (1.3-17.2)	98 (88.9-99.95)
	N	48	48
	hSBA GMT (95% CI)	1.17 (1.00-1.37)	45.09 (35 - 59)
	N	48	48
NadA	% seropositive (95% CI)	7 (1.4-18.7)	100 (92.3-100)
	N	44	46
	hSBA GMT (95% CI)	1.28 (1.00-1.66)	233.99 (177 - 309)
	N	44	46
PorA P1.4	% seropositive (95% CI)	0 (0-7.4)	77 (62.7-88)
	N	48	48
	hSBA GMT (95% CI)	1.00 (1.00-1.00)	12.39 (8.33 - 18)
	N	48	48
NHBA	% seropositive (95% CI)	54 (39.2-68.6)	88 (74.8-95.3)
	N	48	48
	hSBA GMT (95% CI)	8.97 (5.21-15)	30.38 (19-49)
	N	48	48

Abbreviations: CI = confidence interval; hSBA = human serum bactericidal activity; GMT = geometric mean titers.

<sup>a</sup> Immunogenicity subset. % seropositive = the percentage of subjects who achieved an hSBA  $\geq$  1:5.

<sup>b</sup> N=145 follow on subjects and N=105 for naïve subjects

N= number of subjects with evaluable serum for analysis of that strain.

In study V72\_75, antibody persistence data were obtained in Canada and Australia (V72\_41), and in Chile (V72P10). At approximately 4 years, hSBA GMTs declined after the 2-dose primary series as adolescents in Canada and Australia. The percentages of subjects with hSBA  $\geq$  1:5 were 26% for fHbp, 84% for 5/99NadA, 9% for PorA P1.4, and 71% for NHBA. The response to a booster dose after 4 years was indicative of immunological memory as 98% of subjects reached an hSBA  $\geq$  1:5 to fHbp, 100% to NadA, 92% to PorA P1.4, and 99% to NHBA, respectively. At approximately 7.5 years, hSBA GMTs declined after the 2-dose primary series as adolescent in Chile. The percentage of subjects with hSBA  $\geq$  1:4 was 44% for fHbp, 84% for NadA, 29% for PorA P1.4, and 78% for NHBA. The response to a booster dose after 7.5 years was indicative of immunological memory as 100% of subjects reached an hSBA  $\geq$  1:4 to fHbp, 100% to NadA, 93% to PorA P1.4, and 98% to NHBA, respectively.

Limited immunogenicity data are available in subjects above 25 years of age in studies V72P4, V72P5 and V72P59. Ranges for the percentages of subjects with pre-existing hSBA  $\geq$  1:4/1:5 across studies in subjects above 25 years of age were: 0% to 39% for fHbp, 33% to 41% for NadA, 0% to 22% for PorA P.4. All subjects (100%) in V72\_59 had hSBA  $\geq$  1:5 for NHBA. At 1 month after a 2-dose schedule of BEXSERO, majority of subjects achieved an hSBA  $\geq$  1:4/1:5 for each of the strains tested (range across studies: 100% for fHbp, 100% for NadA, 73% to 96%

for PorA P1.4; and 100% for NHBA (study V72\_59 only)).

## TOXICOLOGY

**Table 23 - Nonclinical Toxicology Studies**

Study type, gender, and species	Route and regimen	Results
Single and repeat dose toxicity and local tolerability, male and female rabbits	One or five 0.5 mL or 1 mL intramuscular doses of rMenB±OMV <sup>d</sup> (50 mcg or 100 mcg of each recombinant protein NHBA, NadA and fHbp <sup>a</sup> , and 25 mcg of OMV NZ or NW in 1.5 mg or 3 mg Al(OH) <sub>3</sub> ) two weeks apart for eight weeks	No systemic adverse effects and well tolerated locally
Pilot reproductive & developmental toxicity female rabbits	Five 0.5 mL or 1 mL intramuscular doses of rMenB±OMV <sup>d</sup> approx. two weeks apart. Three doses before mating and two during gestation (1× dose in 0.5 mL: 50 mcg of each recombinant protein NHBA, NadA and fHbp <sup>a</sup> , 25 mcg of OMV NZ in 1.5 mg Al(OH) <sub>3</sub> ; or 2× dose in 1 mL, administered 0.5 mL in each leg)	No systemic toxicity in maternal rabbits and no teratogenic effects
Pivotal reproductive & developmental toxicity female rabbits	Five 0.5 mL intramuscular doses of rMenB±OMV <sup>d</sup> two weeks apart. Three doses before mating and two during gestation (50 mcg of each recombinant protein NHBA, NadA and fHbp <sup>a</sup> , 25 mcg of OMV NZ in 1.5 mg Al(OH) <sub>3</sub> )	No systemic toxicity in maternal rabbits and no reproductive, embryofetal, or postnatal developmental effects
In vitro toxicity non-GLP studies in human cells	HBMEC, HUVEC cells <sup>b</sup> , human plasma or whole blood and platelet-rich plasma treated with vaccine components at various incubation times	No effects on cytotoxicity, binding to human cells, cytokines production, coagulation <sup>c</sup> , platelet activation, platelet-leukocyte aggregation
Single and repeat dose toxicity and local tolerability male and female rabbits	One or five 0.5 mL intramuscular doses of MenB protein 287±OMV (50 mcg MenB recombinant 287 ± 25 mcg OMV in 1.65 mg Al(OH) <sub>3</sub> ) two weeks apart	No systemic adverse effects and well tolerated locally
Reproductive & developmental toxicity female rabbits	Eight 0.5 mL intramuscular doses of MeNZB <sup>TM</sup> . Three doses two weeks apart before mating and five doses every 3 to 4 days during gestation (25 mcg OMV in 0.5 mL with 1.65 mg Al(OH) <sub>3</sub> before mating; 6.25 mcg, 25 mcg or 50 mcg OMV in 0.13 to 1 mL with Al(OH) <sub>3</sub> during gestation)	No systemic toxicity in maternal rabbits and no teratogenic effects

<sup>a</sup> proteins NHBA, NadA and fHbp also named antigens 287-953, 961c, and 936-741;

<sup>b</sup> human umbilical vein endothelial cells and human brain microvascular endothelial cells;

<sup>c</sup> PT, PTT and activated Protein C;

PT: prothrombin time;

PTT: partial thromboplastin time;

<sup>d</sup> rMenB+OMV NZ corresponds to BEXSERO

## REFERENCES

1. Bettinger JA, Scheifele DW, Le Saux N, Halperin SA, Vaudry W, Tsang R. The Disease Burden of Invasive Meningococcal Serogroup B Disease in Canada. *Pediatr Infect Dis J* 2013;32: e20–e25.
2. Gilca R *et al.* The Changing Epidemiology of Meningococcal Disease in Quebec, Canada, 1991–2011: Potential Implications of Emergence of New Strains. *PLOS One* 7(11): 2012;7:1–9.

3. Donnelly, Medini D, Boccadifuoco G, Biolchi A, Ward J et al. Qualitative and quantitative assessment of meningococcal antigens to evaluate the potential strain coverage of protein-based vaccines. *Proc Natl Acad Sci U S A* 2010; 107(45):19490-5
4. Julie A. Bettinger, David W. Scheifele, Scott A. Halperin, Wendy Vaudry, Jamie Findlow, Ray Borrow, Duccio Medini, Raymond Tsang, For the members of the Canadian Immunization Monitoring Program, Active (IMPACT), Diversity of Canadian meningococcal serogroup B isolates and estimated coverage by an investigational meningococcal serogroup B vaccine (4CMenB), Vaccine, Available online 12 April 2013, ISSN 0264-410X, <http://dx.doi.org/10.1016/j.vaccine.2013.03.063>.
5. Goldschneider I., et. al. (1969b) Human immunity to the meningococcus. II. Development of natural immunity. *J Exp Med* (6):1327-1348.
6. Goldschneider I., et. al. (1969a) Human immunity to the meningococcus. I. The role of humoral antibodies. *J Exp Med* (6):1307-26.
7. Sadarangani, M., et al. (2015). "Outcomes of invasive meningococcal disease in adults and children in Canada between 2002 and 2011: a prospective cohort study." *Clin Infect Dis* 60(8): e27-35.
8. Kimura A, Toneatto D, Kleinschmidt A, Wang H, Dull P. Immunogenicity and Safety of a Multicomponent Meningococcal Serogroup B Vaccine and a Quadrivalent Meningococcal CRM<sub>197</sub> Conjugate Vaccine against Serogroups A, C, W-135, and Y in Adults Who Are at Increased Risk for Occupational Exposure to Meningococcal Isolates. *Clinical and Vaccine Immunology* 18:3 (2011) 483–486.
9. Toneatto D, Ismaili S, Ypma E, Vienken K, Oster P, Dull P. The first use of an investigational multicomponent meningococcal serogroup B vaccine (4CMenB) in humans. *Human Vaccines* 7:6 (2011) 646-653.
10. Read RC, Dull P, Bai X, Nolan K, Findlow J, Bazaz R, Kleinschmidt A, McCarthy M, Wang H, Toneatto D, Borrow R. A phase III observer-blind randomized, controlled study to evaluate the immune response and the correlation with nasopharyngeal carriage after immunization of university students with a quadrivalent meningococcal ACWY glycoconjugate or serogroup B meningococcal vaccine. *Vaccine* 35 (2017) 427-434.
11. INSPQ (October 2018). Epidemiological impact of the meningococcal serogroup B vaccination campaign in the Saguenay-Lac-Saint-Jean region, 2014: report as of June 30, 2018. Available: <https://www.inspq.qc.ca/publications/2491>.
12. Madico G *et al.* The Meningococcal Vaccine Candidate GNA1870 Binds the Complement Regulatory Protein Factor H and Enhances Serum Persistence. *J Immunol* 2006;177:501–510.
13. Schneider MC *et al.* *Neisseria meningitidis* recruits factor H using protein mimicry of host carbohydrates. *Nature* 2009;458:890–893.

14. Serruto D *et al.* *Neisseria meningitidis* GNA2132, a heparin-binding protein that induces protective immunity in humans. *Proc Natl Acad Sci U S A* 2010;107:3770–3775.
15. Bambini S *et al.* Distribution and genetic variability of three vaccine components in a panel of strains representative of the diversity of serogroup B meningococcus. *Vaccine* 2009;27:2794–2803.
16. Martin DR *et al.* The VR2 Epitope on the PorA P1.7-2,4 Protein Is the Major Target for the Immune Response Elicited by the Strain-Specific Group B Meningococcal Vaccine MeNZB. *Clin Vaccine Immunol* 2006;13:486–491.
17. Vesikari T *et al.* Immunogenicity and safety of an investigational multicomponent, recombinant, meningococcal serogroup B vaccine (4CMenB) administered concomitantly with routine infant and child vaccinations: results of two randomised trials. *Lancet* 2013;381:825–835.
18. Vogel U *et al.* Predicted strain coverage of a meningococcal multicomponent vaccine (4CMenB) in Europe: a qualitative and quantitative assessment. *Lancet Infect Dis* 2013;13:416–425.
19. Capecchi B, Adu-Bobie J, Di Marcello F, *et al.* *Neisseria meningitidis* NadA is a new invasin which promotes bacterial adhesion to and penetration into human epithelial cells. *Mol Microbiol* 2005; 55(3): 687-98.
20. Wang X *et al.* Prevalence and genetic diversity of candidate vaccine antigens among invasive *Neisseria meningitidis* isolates in the United States. *Vaccine* 2011; 29(29-30):4739-44.
21. Comanducci M *et al.* NadA, a novel vaccine candidate of *Neisseria meningitidis*. *J Exp Med* 2002; 195: 1445-1454.
22. Mazzon C *et al.* IFN- $\gamma$  and R-848 Dependent Activation of Human Monocyte-Derived Dendritic Cells by *Neisseria meningitidis* Adhesin A. *J Immunol* 2007; 179: 3904-3916.
23. De Wals P *et al.* Impact of an Immunization Campaign to Control an Increased Incidence of Serogroup B Meningococcal Disease in One Region of Quebec, Canada. *Clin Infect Dis* 2017 May 1; 64(9):1263-1267.



### PART III: CONSUMER INFORMATION

#### BEXSERO

Multicomponent Meningococcal B Vaccine  
(recombinant, adsorbed)

This leaflet is Part III of a three-part "Product Monograph" published when BEXSERO was approved for sale in Canada and designed specifically for consumers. This leaflet is a summary and will not tell you everything about BEXSERO. Contact your doctor/pharmacist/nurse if you have any questions about the vaccine.

#### ABOUT THIS VACCINE

##### What the vaccine is used for:

BEXSERO is a vaccine for the prevention of Meningococcal disease caused by the *Neisseria meningitidis* group B bacteria (germs). These germs can cause serious, and sometimes life-threatening, infections such as meningitis (infection of the lining of the brain and spinal cord) and sepsis (blood poisoning).

BEXSERO is given to individuals from 2 months through 25 years of age.

##### What it does:

The vaccine works by specifically stimulating the immune system of the vaccinated person, causing the production of substances in the blood called antibodies. The antibodies kill the germ that causes meningococcal disease, *N. meningitidis*. If a vaccinated person is infected by *N. meningitidis*, their immune system is usually ready to destroy it.

##### When it should not be used:

If you or your child are allergic (hypersensitive) to the active substances or any of the other ingredients of BEXSERO

##### What the medicinal ingredients are:

The active substances are:

50 mcg of recombinant *Neisseria meningitidis* group B NHBA fusion protein  
50 mcg of recombinant *Neisseria meningitidis* group B NadA protein  
50 mcg of recombinant *Neisseria meningitidis* group B fHbp fusion protein  
25 mcg of Outer Membrane Vesicles *Neisseria meningitidis* group B strain NZ98/254

Antigens are adsorbed on aluminium hydroxide (0.5 mg aluminium).

(mcg = micrograms)

##### What the nonmedicinal ingredients are:

Aluminium hydroxide, histidine, sodium chloride, sucrose, water for injections. Residue from the manufacturing process: kanamycin.

##### What dosage forms are available?

Each dose of 0.5 mL is a suspension for intramuscular injection provided in a prefilled glass (Type I) syringe. Syringes are available in packages containing either one or ten syringes, supplied with or without needles.

#### WARNINGS AND PRECAUTIONS

BEFORE you or your child receive BEXSERO, talk to your doctor/pharmacist/nurse if:

- you or your child have a severe infection with a high temperature. If this is the case, then vaccination will be postponed. The presence of a minor infection, such as a cold, should not require postponement of the vaccination, but talk to your doctor/pharmacist/nurse first.
- If you receive treatment that blocks the part of the immune system known as complement activation, such as eculizumab. Even if you have been vaccinated with BEXSERO you remain at increased risk of disease caused by the *Neisseria meningitidis* group B bacteria.
- you are pregnant or breast feeding, think you may be pregnant or are planning to have a baby, ask your doctor/pharmacist/nurse for advice before BEXSERO is given.
- you have hemophilia or any other condition that may slow down the clotting of your blood, such as treatment with blood thinners (anticoagulants).
- your child was born prematurely (before or at 28 weeks of pregnancy), particularly with breathing difficulties. Stopping breathing or irregular breathing for a short time may be more common in the first three days following vaccination in these babies and they may need special monitoring.
- you or your child have an allergy to the antibiotic kanamycin. If present, the kanamycin level in the vaccine is low. If you or your child may have allergy to kanamycin, talk to your doctor/pharmacist/nurse first.

Fainting, feeling faint or other stress-related reactions can occur as a response to any needle injection. Tell your doctor or nurse if you have experienced this kind of reaction previously.

Tell your doctor/pharmacist/nurse if you know that you or your child is allergic to latex. The tip cap of the syringe may contain natural rubber latex. Although the risk for developing allergic reactions is very small, your doctor/pharmacist/nurse should consider the benefit-risk prior to administering this vaccine to individuals with known history of hypersensitivity to latex.

Your doctor/pharmacist/nurse may ask you to give your child medicines that lower fever at the time and after BEXSERO has been given. This will help to reduce some of the side effects of BEXSERO.

There are limited data on the use of BEXSERO in patients with chronic medical conditions or with weakened immunity. If you or your child have weakened immunity (for example, due to the use of immunosuppressive medications, or HIV infection, or hereditary defects of the body's natural defense system), it is possible that the effectiveness of BEXSERO is reduced.

As with any vaccine, BEXSERO may not fully protect all of those who are vaccinated.

BEXSERO is not expected to provide protection against all circulating meningococcal serogroup B strains.

**INTERACTIONS WITH THIS VACCINE**

Tell your doctor/pharmacist/nurse if you or your child are taking, have recently taken, or might take any other medicines, or have recently received any other vaccine.

BEXSERO can be given at the same time as any of the following vaccine antigens, either as monovalent or as combination vaccines: diphtheria, tetanus, acellular and pertussis (whooping cough), *Haemophilus influenzae* type b, inactivated polio, hepatitis B, heptavalent pneumococcal conjugate, measles, mumps, rubella, chickenpox, and meningococcal groups A, C, W, Y conjugate. Talk to your doctor/pharmacist/nurse for further information.

When BEXSERO is given at the same time as any other vaccine, the vaccines must be given at separate sites.

BEXSERO has no or negligible influence on the ability to drive and use machines. However, some of the effects mentioned under section "Side effects and what to do about them" may temporarily affect the ability to drive or use machines.

**PROPER USE OF THIS VACCINE**

Usual dose:

Your doctor/pharmacist/nurse will inject the recommended dose (0.5 mL) of the vaccine into your or your child's arm or leg muscle.

BEXSERO must not be mixed with any other vaccine or medicinal products in the same syringe.

***Infants aged 2 months to 5 months at the time of first dose***

Your child should receive an initial course of two or three injections of the vaccine followed by an additional injection (booster).

The interval between vaccinations should be at least 2 months if two initial doses are given or at least 1 month if three initial doses are given. A booster will be given in the second year of life after an interval of at least 6 months from the last injection of the initial course.

***Infants aged 6 months to 11 months of age at the time of first dose***

Your child should receive two injections of the vaccine, given at least 2 months apart. A booster will be given in the second year of life, after an interval of at least 2 months from the last dose.

***Children aged 12 months to 23 months at the time of first dose***

Your child should receive two injections of the vaccine, given at least 2 months apart. A booster will be given after an interval of 12 to 23 months from the second injection.

***Children, Adolescents and Adults aged 2 through 25 years at the time of first dose***

You or your child should receive two injections, given at least 1 month apart. An additional booster injection may be considered in individuals at continuous risk of exposure to meningococcal disease.

Make sure that you or your child gets all doses. This allows you or your child to get the full benefits of BEXSERO.

Missed Dose:

If you forget to go back to the doctor/pharmacist/nurse at the scheduled time ask the doctor/pharmacist/nurse for advice. If you have any further questions on the use of BEXSERO, ask your doctor/pharmacist/nurse.

Overdose:

In case of vaccine overdose, contact a health care practitioner, hospital emergency department or regional Poison Control Centre immediately, even if there are no symptoms.

**SIDE EFFECTS AND WHAT TO DO ABOUT THEM**

Like all vaccines, BEXSERO can cause side effects, although not everybody gets them.

When BEXSERO is given to you or your child, the very common side effects (may affect more than 1 in 10 people) that you or your child may get (reported in all age groups) are:

- pain/tenderness, redness, swelling, or hardness of the skin at the injection site.

The following side effects may also occur after receiving this vaccine.

**Infants and children (2 months to 10 years of age)**

Very common (these may affect more than 1 in 10 people)

- fever ( $\geq 38^{\circ}\text{C}$ )
- loss of appetite
- tenderness at the injection site (including severe injection site tenderness resulting in crying when injected limb is moved)
- skin rash (uncommon after booster)
- sleepiness
- feeling irritable
- unusual crying
- vomiting (uncommon after booster)
- diarrhea
- headache
- painful joints

Uncommon (these may affect up to 1 in 100 people)

- high fever ( $\geq 40^{\circ}\text{C}$ )
- seizures (including febrile seizures)
- dry skin, itchy rash, skin rash
- paleness (rare after booster)
- itchy rash, skin rash

Rare (these may affect up to 1 in 1,000 people)

- Kawasaki disease which may include symptoms such as fever that lasts for more than five days, associated with a skin rash on the trunk of the body, and sometimes followed by a peeling of the skin on the hands and fingers, swollen glands in the neck, red eyes, lips, throat and tongue.

#### Adolescents and Adults (11 years of age and older)

Very common (these may affect more than 1 in 10 people).

- pain at the injection site resulting in inability to perform normal daily activity
- painful muscles and joints
- nausea
- generally feeling unwell
- headache

Side effects that have been reported during marketed use include:

- Allergic reactions that may include severe swelling of the lips, mouth, throat (which may cause difficulty in swallowing), difficulty breathing with wheezing or coughing, rash, loss of consciousness and very low blood pressure;
- collapse (sudden onset of muscle floppiness), less responsive than usual or lack of awareness, and paleness or bluish skin discoloration in young children; feeling faint or fainting;
- fever (adolescents from 11 years of age and adults); injection site reactions like extensive swelling of the vaccinated limb, blisters at or around the injection site and hard lump at the injection site (which may persist for more than one month);
- skin rash (adolescents from 11 years of age and adults).

If any of the noted side effects becomes serious, or if you notice any side effects not listed in this leaflet, please tell your doctor/pharmacist/nurse immediately.

*This is not a complete list of side effects. For any unexpected effects while taking BEXSERO, contact your doctor / pharmacist / nurse.*

#### Reporting Suspected Side Effects

**For the general public:** Should you experience a side effect following immunization, please report it to your doctor, nurse, or pharmacist.

Should you require information related to the management of the side effect, please contact your healthcare provider. The Public Health Agency of Canada, Health Canada and GlaxoSmithKline Inc. cannot provide medical advice.

**For healthcare professionals:** If a patient experiences a side effect following immunization, please complete the Adverse Events Following Immunization (AEFI) Form appropriate for your province/territory (<http://www.phac-aspc.gc.ca/im/aeffi-esssi-form-eng.php>) and send it to your local Health Unit

#### HOW TO STORE IT

Store in a refrigerator at  $2^{\circ}\text{C}$  to  $8^{\circ}\text{C}$ . Do not freeze. Do not use vaccine that may have been frozen. Protect from light. Do not use BEXSERO after the expiry date. Keep out of reach of children.

#### MORE INFORMATION

This document plus the full product monograph, prepared for health professionals can be found at: [www.gsk.ca](http://www.gsk.ca) or by contacting the sponsor, GlaxoSmithKline Inc. 7333 Mississauga Road, Mississauga, Ontario L5N 6L4 1-800-387-7374

This leaflet was prepared by GlaxoSmithKline Inc.  
Last revised: March 11, 2020

©2020 GSK group of companies or its licensor  
Trademarks are owned by or licensed to the GSK group of companies.